

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
8 April 2004 (08.04.2004)

PCT

(10) International Publication Number  
**WO 2004/029616 A2**

(51) International Patent Classification<sup>7</sup>: G01N 33/48

(21) International Application Number:  
PCT/CA2003/001479

(22) International Filing Date:  
26 September 2003 (26.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/414,314 26 September 2002 (26.09.2002) US

(71) Applicant (for all designated States except US): MOUNT SINAI HOSPITAL [CA/CA]; 600 University Avenue, Toronto, Ontario M5G 1X5 (CA).

(72) Inventor; and

(75) Inventor/Applicant (for US only): DIAMANDIS, Eleftherios, P. [CA/CA]; 44 Gerrard Street West, Suite 1504, Toronto, Ontario M5G 2K2 (CA).


(74) Agent: KURDYDYK, Linda; McCarthy Tetrault, Suite 4700, P.O. Box 48, 66 Wellington Street West, Toronto Dominion Bank Tower, Toronto, Ontario M5K 1E6 (CA).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: METHODS FOR DETECTING PROSTATE CANCER



(57) Abstract: The invention relates to the application of kallikrein (11), free PSA, and total PSA in the detection of prostate cancer. These markers may be used for the diagnosis, monitoring, staging, progression, prevention, treatment, and prognosis of prostate cancer, and as indicators before surgery or after relapse. A particular aspect of the invention provides a method for distinguishing prostate cancer from benign prostatic hyperplasia (BPH).



**Published:**

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

- 1 -

**TITLE: METHODS FOR DETECTING PROSTATE CANCER****FIELD OF THE INVENTION**

The invention relates to kallikrein 11, prostate-specific antigen, free prostate-specific antigen and their use in detecting prostate cancer.

**5 BACKGROUND OF THE INVENTION**

Prostate cancer (CaP) is the most frequently diagnosed cancer in men in North America and its mortality rate is second only to lung cancer. Therefore, early diagnosis and monitoring of prostate cancer is an important priority. Prostate-specific antigen (PSA) is widely used as the most reliable tumor marker established so far<sup>1,2</sup>. However, non-malignant prostatic diseases, especially benign prostatic hyperplasia (BPH) and acute prostatitis also cause serum PSA elevation, thus complicating the diagnosis of prostate cancer by PSA measurements alone. Analysis of the molecular forms of PSA improves specificity for prostate cancer<sup>3,4</sup>. The determination of free PSA and its ratio to total PSA is now clinically established and is used to reduce the number of unnecessary prostate biopsies<sup>5</sup>. Despite numerous efforts to further reduce unnecessary biopsies, false negative and false positive results still occur with high frequency.

15 The human kallikrein 11 gene (KLK11) was originally isolated from human hippocampus as trypsin-like serine protease (TLSP)<sup>6</sup>. Two alternative splice variants of this gene, also known as hippostasin have also been identified<sup>7-9</sup>. With the official nomenclature, TLSP/hippostasin is now known as human kallikrein 11 (hK11). This protein is encoded by the KLK11 gene that belongs to the human kallikrein family along with PSA (hK3) and other kallikreins<sup>10</sup>. hK11 protein has been found to be highly expressed in the prostate<sup>8,11</sup>.

The citation of any reference herein is not an admission that such reference is available as prior art to the instant invention.

**SUMMARY OF THE INVENTION**

The invention relates to the application of kallikrein 11, free PSA, and total PSA in the detection of prostate cancer. These markers may be used for the diagnosis, monitoring, staging, progression, prevention, treatment, and prognosis of prostate cancer, and as indicators before surgery or after relapse.

Kallikrein 11, and PSA, and agents that bind to kallikrein 11 and PSA, may be used to detect prostate cancer and they can be used in the diagnostic evaluation of prostate cancer, and the identification of subjects with a predisposition to such disorders. Methods for detecting kallikrein 11 and PSA can be used to monitor prostate cancer.

The presence of kallikrein 11 and PSA in a sample can be assessed, for example, by detecting the presence in the sample of (a) kallikrein 11 and PSA or fragments thereof; or (b) metabolites which are produced directly or indirectly by a kallikrein 11 or PSA.

In an aspect, the invention provides a method for detecting prostate cancer in a subject comprising measuring kallikrein 11 and prostate specific antigen (PSA) in a sample from the subject.

In an embodiment, the invention provides a method for detecting kallikrein 11 and PSA comprising (a) obtaining a sample from a patient; (b) detecting or identifying in the sample kallikrein 11 and PSA; and (c) comparing the detected amounts with amounts detected for a standard.

In an aspect of the invention, a method for screening a subject for prostate cancer is provided

- 2 -

comprising (a) obtaining a biological sample from a subject; (b) detecting the amounts of kallikrein 11, free PSA, and total PSA in said sample; and (c) comparing said amounts of kallikrein 11, free PSA, and total PSA detected to predetermined standards, where detection of different levels of kallikrein 11, free PSA, and total PSA compared with that of a standard indicates disease.

5       The terms "detect", "detecting" and "detection" include assaying or otherwise establishing the presence or absence of kallikrein 11, a combination of kallikrein 11 and PSA, subunits thereof, or combinations of reagent bound targets, and the like, or assaying for, imaging, ascertaining, establishing, or otherwise determining one or more factual characteristics of prostate cancer, metastasis, stage, or similar conditions. The term encompasses diagnostic, prognostic, and monitoring applications.

10       According to a method involving kallikrein 11 and PSA, the levels in a sample from a patient are compared with the normal levels of kallikrein 11 and PSA in samples of the same type obtained from controls (e.g. samples from individuals not afflicted with disease). Significantly altered levels in the sample of kallikrein 11 and PSA relative to the normal levels in a control is indicative of disease. Thus, a method is provided of assessing whether a patient is afflicted with or has a pre-disposition to prostate cancer,  
15       comprising comparing (a) levels of kallikrein 11 and PSA in a sample from the patient; and (b) normal levels of kallikrein 11 and PSA in samples of the same type obtained from control patients not afflicted with prostate cancer, wherein significantly altered levels of kallikrein 11 and PSA relative to the corresponding normal levels of kallikrein 11 and PSA, is an indication that the patient is afflicted with or has a pre-disposition to prostate cancer.

20       A significant difference between the levels of kallikrein 11 and PSA in the patient and the normal levels (e.g. lower levels in the patient) is an indication that the patient is afflicted with or has a predisposition to prostate cancer.

      The present invention also relates to a method for diagnosing and monitoring prostate cancer in a subject comprising detecting or measuring kallikrein 11 and PSA in a sample from the subject using a  
25       reagent that detects kallikrein 11 and PSA, in particular antibodies specifically reactive with kallikrein 11 and PSA or a part thereof. In an embodiment, the sample is serum.

      In an embodiment, the invention relates to a method for diagnosing and monitoring prostate cancer in a subject by quantitating kallikrein 11 and PSA in a biological sample from the subject comprising (a) reacting the biological sample with antibodies specific for kallikrein 11 and PSA which is directly or  
30       indirectly labelled with a detectable substance; and (b) detecting the detectable substance.

      In another aspect the invention provides a method for using antibodies to detect expression of a kallikrein 11 and PSA in a sample, the method comprising: (a) combining antibodies specific for kallikrein 11 and PSA with a sample under conditions which allow the formation of antibody:protein complexes; and (b) detecting complex formation, wherein complex formation indicates expression of kallikrein 11 and PSA  
35       in the sample. Expression may be compared with standards and is diagnostic of prostate cancer.

      Embodiments of the methods of the invention involve (a) reacting a biological sample from a subject with antibodies specific for kallikrein 11 and PSA which are directly or indirectly labelled with an enzyme; (b) adding a substrate for the enzyme wherein the substrate is selected so that the substrate, or a reaction product of the enzyme and substrate forms fluorescent complexes; (c) quantitating kallikrein 11 and

- 3 -

PSA in the sample by measuring fluorescence of the fluorescent complexes; and (d) comparing the quantitated levels to levels obtained for other samples from the subject patient, or control subjects. In an embodiment the quantitated levels are compared to levels quantitated for control subjects without prostate cancer wherein lower levels of kallikrein 11 and PSA compared with the control subjects is indicative of prostate cancer.

A particular embodiment of the invention comprises the following steps

- (a) incubating a biological sample with first antibodies specific for kallikrein 11 and PSA which are directly or indirectly labeled with detectable substances, and second antibodies specific for kallikrein 11 and PSA which are immobilized;
- (b) detecting the detectable substances thereby quantitating kallikrein 11 and PSA in the biological sample; and
- (c) comparing the quantitated kallikrein 11 and PSA with levels for a predetermined standard.

The standard may correspond to levels quantitated for samples from control subjects without prostate cancer, with a different disease stage, or from other samples of the subject. Lower levels of kallikrein 11 and PSA as compared to the standard is indicative of prostate cancer.

The invention also contemplates the methods described herein using multiple markers for prostate cancer. Therefore, the invention contemplates a method for analyzing a biological sample for the presence of kallikrein 11 and PSA and other markers that are specific indicators of prostate cancer. The methods described herein may be modified by including reagents to detect the markers, or nucleic acids for the markers.

The invention further relates to a method of assessing the efficacy of a therapy for inhibiting prostate cancer in a patient. A method of the invention comprises comparing: (a) levels of kallikrein 11 and PSA in a sample from the patient obtained from the patient prior to providing at least a portion of the therapy to the patient; and (b) levels of kallikrein 11 and PSA in a second sample obtained from the patient following therapy.

A significant difference between the levels of kallikrein 11 and PSA in the second sample relative to the first sample (e.g. higher levels of kallikrein 11 and PSA) is an indication that the therapy is efficacious for inhibiting prostate cancer.

The "therapy" may be any therapy for treating prostate cancer including but not limited to therapeutics, radiation, immunotherapy, gene therapy, and surgical removal of tissue. Therefore, the method can be used to evaluate a patient before, during, and after therapy.

In an aspect, the invention provides a method for monitoring the progression of prostate cancer in a patient the method comprising:

- (a) detecting kallikrein 11 and PSA in a sample from the patient at a first time point;
- (b) repeating step (a) at a subsequent point in time; and
- (c) comparing the levels detected in (a) and (b), and therefrom monitoring the progression of the prostate cancer.

The invention also provides a method for assessing the potential efficacy of a test agent for inhibiting prostate cancer, and a method of selecting an agent for inhibiting prostate cancer.

- 4 -

The invention also contemplates a method of assessing the potential of a test compound to contribute to prostate cancer comprising:

- (a) maintaining separate aliquots of prostate cancer diseased cells in the presence and absence of the test compound; and
- 5 (b) comparing the levels of kallikrein 11 and PSA in each of the aliquots.

A significant difference between the levels of kallikrein 11 and PSA in the aliquot maintained in the presence of (or exposed to) the test compound relative to the aliquot maintained in the absence of the test compound, indicates that the test compound potentially contributes to prostate cancer.

In the methods of the invention the amount of kallikrein 11 and the amount of PSA may be mathematically combined. In an embodiment of the invention, a method is provided for detecting or  
10 identifying prostate cancer in a subject comprising:

- (a) determining the amount of kallikrein 11 in a sample from the subject;
- (b) determining the amount of PSA in the sample;
- (c) mathematically combining the results of step (a) and step (b); and
- 15 (d) relating the combination to the presence of prostate cancer.

In aspects of the invention, the combination is a ratio of kallikrein 11 to total PSA, or the inverse thereof. The combination is preferably compared to a mathematical combination for a predetermined standard.

In particular, the invention provides a method for screening or identifying prostate cancer by  
20 determining the ratio between kallikrein 11:total PSA in a sample from a subject, preferably a serum sample.

A method of the invention, in particular a method for detecting or identifying prostate cancer, may further comprise determining the % free PSA and relating the combination and % free PSA to the presence of prostate cancer.

25 In an aspect the invention provides a method for distinguishing prostate cancer from benign prostatic hyperplasia (BPH) in a subject comprising determining the amount of kallikrein 11 contained in a sample from the subject, and relating the amount to the presence of prostate cancer or BPH in the subject. The amount of kallikrein 11 in the sample may be compared to an amount determined for a standard. A standard may be an amount of kallikrein 11 associated with prostate cancer, and a higher amount of  
30 kallikrein 11 in the sample compared to the standard may be indicative of BPH. A standard may be an amount of kallikrein 11 associated with BPH, and a lower amount of kallikrein 11 in the sample compared to the standard may be indicative of prostate cancer.

In a further aspect, the invention provides a method for distinguishing prostate cancer from benign prostatic hyperplasia (BPH) in a subject comprising:

- 35 (a) determining the amount of kallikrein 11 contained in a sample from the subject;
- (b) determining the amount of total PSA contained in the sample;
- (c) mathematically combining the results of (a) and (b);
- (d) relating the combination to the presence of BPH or prostate cancer.

The combination may be a ratio of kallikrein 11 to total PSA, or the inverse thereof.

- 5 -

Therefore, the invention provides a method for differentiation between BPH or prostate cancer by determining the ratio between kallikrein 11:total PSA in a subject's serum.

The combination (e.g. ratio) may be compared to a combination for a standard associated with prostate cancer or BPH. A standard may be a combination (e.g. ratio) determined for prostate cancer, and a  
5 higher combination (e.g. ratio) determined for the sample compared to the standard may be indicative of BPH. A standard may be a combination (e.g. ratio) determined for BPH, and a lower combination determined for the sample compared to the standard may be indicative of prostate cancer.

The methods of the invention may further comprise determining the percentage of free PSA and correlating the percentage free PSA and the combination to the presence of prostate cancer or BPH in the  
10 subject.

In the methods of the invention kallikrein 11 and PSA may be measured using agents that bind kallikrein 11 and PSA, respectively. The binding agents may be directly or indirectly labeled with a detectable substance.

In particular, the invention provides a diagnostic method for determining the presence of BPH or  
15 prostate cancer in a subject comprising:

- (a) contacting a sample from the subject with a first binding agent that binds to kallikrein 11 and a second binding agent that binds to PSA;
- (b) determining the presence or amount of first complexes comprising the first agent and kallikrein 11 and second complexes comprising the second agent and PSA;
- 20 (c) mathematically combining the amount of the first and second complexes; and
- (d) relating the combination to the presence of prostate cancer.

In particular methods of the invention, patients with BPH or prostate cancer can be identified in patients with total PSA between about 4 to 10 ng/ml.

In still other particular methods of the invention, patients with BPH or prostate cancer can be  
25 identified in patients with total PSA less than 4 ng/ml.

The invention also provides a method of improving the accuracy of a diagnosis of prostate cancer comprising the steps of: a) performing a method of the invention; and b) performing at least one of a test for free PSA and a digital rectal examination.

The invention also relates to kits for carrying out the methods of the invention. In an embodiment,  
30 the kit is for assessing whether a patient is afflicted with prostate cancer, and it comprises reagents for assessing kallikrein 11 and PSA.

In another aspect, the invention relates to a kit for assessing the suitability of each of a plurality of test compounds for inhibiting prostate cancer in a patient. The kit comprises reagents for assessing kallikrein 11 and PSA and optionally a plurality of test agents or compounds.

35 The invention contemplates a kit for assessing the presence of prostate cancer cells, wherein the kit comprises antibodies specific for kallikrein 11 and PSA, and optionally antibodies specific for other markers associated with prostate cancer.

Additionally the invention provides a kit for assessing the potential of a test compound to contribute to prostate cancer. The kit comprises prostate cancer cells and reagents for assessing kallikrein 11 and PSA,

- 6 -

and optionally other markers associated with prostate cancer.

The invention also provides a diagnostic composition comprising kallikrein 11 and PSA or agents that bind to kallikrein 11 and PSA or parts thereof. Agents can be labelled with detectable substances.

The invention also contemplates the methods, compositions, and kits described herein using additional markers associated with prostate cancer. The methods described herein may be modified by including reagents to detect the additional markers, or nucleic acids for the markers.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

#### DESCRIPTION OF THE DRAWINGS

The above-mentioned and other features of this invention and the manner of obtaining them will become more apparent, and will be best understood, by reference to the following description, taken in conjunction with the accompanying drawings. These drawings depict only a typical embodiment of the invention and do not therefore limit its scope. They serve to add specificity and detail, in which:

Figure 1 shows the distribution of hK11/total PSA ratio in BPH and CaP patients. At 90 % sensitivity (hK11/total PSA ratio of 0.06) the specificity is 48.5%. P value was determined by the Mann-Whitney U test.

Figure 2 shows the distribution of hK11/total PSA ratio in the subgroup of patients with % free PSA < 20. At 90 % sensitivity (hK11/total PSA ratio of 0.05) the specificity is 54%. P value was determined by the Mann-Whitney U test.

Figure 3 shows receiver operating characteristic (ROC) curves for total PSA (tPSA), % free PSA and hK11/total PSA ratio, demonstrating the relative potential of each variable in the discrimination of BPH from CaP. AUC, area under the ROC curve; CI, confidence interval

Figure 4 shows the percentage of potentially avoidable biopsies by various biochemical parameters. Eight % of BPH patients would have avoided biopsies by combining of % free PSA and the hK11/tPSA ratio (at 90 % sensitivity).

Figure 5 shows receiver operating characteristic (ROC) curves for total PSA, % free PSA/Total PSA ratio, and hK11/total PSA ratio, demonstrating the relative potential of each variable in the discrimination of BPH from CaP. AUC, area under the ROC curve; CI, confidence interval

Figure 6 shows the distribution of hK11/tPSA ratio in BPH and CaP patients.

Figure 7 shows the distribution of hK11/tPSA ratio in BPH and CaP patients where total PSA = 4-10 µg/L.

Figure 8 shows the distribution of hK11/tPSA ratio in BPH and CaP patients where total PSA = >10 µ/L.

Figure 9 shows the distribution of hK11/tPSA ratio in BPH and CaP patients where total PSA = <4 µ/L.



- 7 -

Figure 10 shows the distribution of hK11/tPSA ratio in BPH and CaP patients where total PSA = < 17.8µ/L (90% sensitivity).

#### **DETAILED DESCRIPTION OF THE INVENTION**

The invention relates to newly discovered correlations between expression of kallikrein 11 in combination with PSA and prostate cancer. The methods described herein provide sensitive methods for detecting prostate cancer. The level of expression of kallikrein 11 and PSA correlates with the presence of prostate cancer. The invention is also based on the finding that kallikrein 11 levels and kallikrein11:PSA ratio are significantly lower in prostate cancer patients than in benign prostatic hyperplasia.

Methods are provided for detecting the presence of prostate cancer in a sample, the absence of prostate cancer in a sample, assessing the histology of tissues associated with prostate cancer, and other characteristics of prostate cancer that are relevant to prevention, diagnosis, characterization, and therapy of prostate cancer in a patient. Methods are also provided for assessing the efficacy of one or more test agents for inhibiting kallikrein 11 and PSA that affect prostate cancer, assessing the efficacy of a therapy for prostate cancer, monitoring the progression of prostate cancer, selecting an agent or therapy for inhibiting prostate cancer, treating a patient afflicted with prostate cancer, inhibiting prostate cancer in a patient, and assessing the potential of a test compound to contribute to prostate cancer.

The invention particularly provides methods and kits for detecting prostate cancer and differentiating prostate cancer and benign prostatic hyperplasia in a subject by determining amounts of kallikrein 11 alone, or kallikrein 11 and PSA in samples from the subjects

#### **Glossary**

Samples that may be analyzed using the methods of the invention include those which are known or suspected to express kallikrein 11 and PSA or contain kallikrein 11 and PSA. The terms "sample", "biological sample", and the like mean a material known or suspected of expressing or containing kallikrein 11 and PSA, in particular kallikrein 11 and PSA associated with prostate cancer. The test sample can be used directly as obtained from the source or following a pretreatment to modify the character of the sample. The sample can be derived from any biological source, such as tissues, extracts, or cell cultures, including cells (e.g. tumor cells), cell lysates, and physiological fluids, such as, for example, whole blood, plasma, serum, saliva, ocular lens fluid, cerebral spinal fluid, sweat, urine, milk, ascites fluid, synovial fluid, peritoneal fluid and the like.

The sample can be obtained from animals, preferably mammals, most preferably humans. The sample can be treated prior to use, such as preparing plasma from blood, diluting viscous fluids, and the like. Methods of treatment can involve filtration, distillation, extraction, concentration, inactivation of interfering components, the addition of reagents, and the like. Proteins may be isolated from the samples and utilized in the methods of the invention.

In embodiments of the invention the sample is a mammalian tissue sample. In another embodiment the sample is a human physiological fluid. In a particular embodiment, the sample is human serum, seminal plasma, urine, or plasma, most preferably serum.

The terms "subject", "individual" or "patient" refer to a warm-blooded animal such as a mammal, which is afflicted with or suspected of having or being pre-disposed to prostate cancer or condition as

described herein. In particular, the terms refer to a human.

The term "kallikrein 11", "kallikrein 11 polypeptide" or "kallikrein 11 protein" includes human kallikrein 11("hK11"), in particular the native-sequence polypeptide, isoforms, chimeric polypeptides, all homologs, fragments, and precursors of human kallikrein 11. The amino acid sequence for native hK11  
5 include the sequences of GenBank Accession Nos. BAA33404 and AB012917 and shown in SEQ ID NO. 1.

A "native-sequence polypeptide" comprises a polypeptide having the same amino acid sequence of a polypeptide derived from nature. Such native-sequence polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term specifically encompasses naturally occurring truncated or secreted forms of a polypeptide, polypeptide variants including naturally occurring variant  
10 forms (e.g. alternatively spliced forms or splice variants), and naturally occurring allelic variants.

The term "polypeptide variant" means a polypeptide having at least about 70-80%, preferably at least about 85%, more preferably at least about 90%, most preferably at least about 95% amino acid sequence identity with a native-sequence polypeptide, in particular having at least 70-80%, 85%, 90%, 95% amino acid sequence identity to the sequences identified in the GenBank Accession Nos. BAA33404 and  
15 AB012917, and shown in SEQ ID NO. 1. Such variants include, for instance, polypeptides wherein one or more amino acid residues are added to, or deleted from, the N- or C-terminus of the full-length or mature sequences of SEQ ID NO: 1, including variants from other species, but excludes a native-sequence polypeptide.

An allelic variant may also be created by introducing substitutions, additions, or deletions into a  
20 nucleic acid encoding a native polypeptide sequence such that one or more amino acid substitutions, additions, or deletions are introduced into the encoded protein. Mutations may be introduced by standard methods, such as site-directed mutagenesis and PCR-mediated mutagenesis. In an embodiment, conservative substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which an amino acid residue is replaced with an amino acid residue with a similar  
25 side chain. Amino acids with similar side chains are known in the art and include amino acids with basic side chains (e.g. Lys, Arg, His), acidic side chains (e.g. Asp, Glu), uncharged polar side chains (e.g. Gly, Asp, Glu, Ser, Thr, Tyr and Cys), nonpolar side chains (e.g. Ala, Val, Leu, Iso, Pro, Trp), beta-branched side chains (e.g. Thr, Val, Iso), and aromatic side chains (e.g. Tyr, Phe, Trp, His). Mutations can also be introduced randomly along part or all of the native sequence, for example, by saturation mutagenesis.  
30 Following mutagenesis the variant polypeptide can be recombinantly expressed and the activity of the polypeptide may be determined.

Polypeptide variants include polypeptides comprising amino acid sequences sufficiently identical to or derived from the amino acid sequence of a native polypeptide which include fewer amino acids than the full length polypeptides. A portion of a polypeptide can be a polypeptide which is for example, 10, 15, 20,  
35 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100 or more amino acids in length. Portions in which regions of a polypeptide are deleted can be prepared by recombinant techniques and can be evaluated for one or more functional activities such as the ability to form antibodies specific for a polypeptide.

A naturally occurring allelic variant may contain conservative amino acid substitutions from the native polypeptide sequence or it may contain a substitution of an amino acid from a corresponding position

in a kallikrein polypeptide homolog, for example, the murine kallikrein polypeptide.

The invention also includes polypeptides that are substantially identical to the sequences of GenBank Accession Nos. BAA33404 and AB012917 and shown in SEQ ID NO. 1 (e.g. at least about 45%, preferably 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, or 99% sequence identity),  
 5 and in particular polypeptides that retain the immunogenic activity of the corresponding native-sequence polypeptide.

Percent identity of two amino acid sequences, or of two nucleic acid sequences identified herein is defined as the percentage of amino acid residues or nucleotides in a candidate sequence that are identical with the amino acid residues in a polypeptide or nucleic acid sequence, after aligning the sequences and  
 10 introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid or nucleic acid sequence identity can be achieved in various conventional ways, for instance, using publicly available computer software including the GCG program package (Devereux J. et al., Nucleic  
 Acids Research 12(1): 387, 1984); BLASTP, BLASTN, and FASTA (Altschul, S.F. et al. J. Molec. Biol. 215:  
 15 403-410, 1990). The BLAST X program is publicly available from NCBI and other sources (BLAST Manual, Altschul, S. et al. NCBI NLM NIH Bethesda, Md. 20894; Altschul, S. et al. J. Mol. Biol. 215: 403-410, 1990). Skilled artisans can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. Methods to determine identity and similarity are codified in publicly available computer programs.

20 Kallikrein 11 proteins include chimeric or fusion proteins. A "chimeric protein" or "fusion protein" comprises all or part (preferably biologically active) of a kallikrein 11 polypeptide operably linked to a heterologous polypeptide (i.e., a polypeptide other than a kallikrein 11 polypeptide). Within the fusion protein, the term "operably linked" is intended to indicate that a kallikrein 11 polypeptide and the heterologous polypeptide are fused in-frame to each other. The heterologous polypeptide can be fused to the  
 25 N-terminus or C-terminus of a kallikrein 11 polypeptide. A useful fusion protein is a GST fusion protein in which a kallikrein 11 polypeptide is fused to the C-terminus of GST sequences. Another example of a fusion protein is an immunoglobulin fusion protein in which all or part of a kallikrein 11 polypeptide is fused to sequences derived from a member of the immunoglobulin protein family. Chimeric and fusion proteins can be produced by standard recombinant DNA techniques.

30 Kallikrein polypeptides may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods, or by any combination of these and similar techniques.

The terms "KLK11" or "KLK11 nucleic acid(s)" are intended to include DNA and RNA (e.g. mRNA) and can be either double stranded or single stranded. The terms include but are not limited to nucleic  
 35 acids that encode a native-sequence polypeptide, a polypeptide variant including a portion of a kallikrein polypeptide, an isoform, precursor, and chimeric polypeptide. The nucleic acid sequences encoding native kallikrein polypeptides employed in the present invention include the nucleic acid sequences of GenBank Accession No. Accession Nos. AF164623 and AB012917 and in SEQ ID NO: 2 and 3, or fragments thereof.

Prostate specific antigen (PSA) is a 33-kDa glycosylated single chain serine protease (Lilja, J Clin

- 10 -

Invest 1985; Watt et al. Proc Natl Acad Sci (USA) 1986). PSA has been predicted to be produced as an inactive zymogen (Lundwall et al. FEBS Lett 1987). Active PSA is secreted into the seminal plasma (Lilja, J Clin Invest 1985) and it is one of the most abundant proteins of the prostate (Lilja et al. The Prostate 1988; Dub et al. J Androl 1987). PSA is capable of forming complexes with serum protease inhibitors. Human  
5 serum contains high levels of  $\alpha_1$ -antichymotrypsin (ACT) and  $\alpha_2$ -macroglobulin, both of which form complexes with PSA. About 70-95% of the PSA in serum that can be detected by immunoassay is complexed with ACT. About 30% of serum PSA does not form a complex with ACT. This non-complexed fraction of PSA contains an internal peptide bond cleavage at Lys145 which renders it inactive. PSA in a complex with a protease inhibitor is referred to "complexed PSA", and PSA that is not complexed with a  
10 protease inhibitor is referred to herein as "free PSA".

The term "total PSA" in reference to a sample refers to the total complexed PSA and free PSA in the sample.

"Statistically different levels" or "significant difference" in levels of markers in a patient sample compared to a control or standard (e.g. normal levels or levels in other samples from a patient) may represent  
15 levels that are higher or lower, in particular lower, than the standard error of the detection assay.

"Binding agent" refers to a substance such as a polypeptide or antibody that specifically binds to a kallikrein 11 or PSA. A substance "specifically binds" to kallikrein 11 or PSA if it reacts at a detectable level with a kallikrein 11 or PSA, and does not react detectably with peptides containing unrelated sequences or sequences of different polypeptides. Binding properties may be assessed using an ELISA, which may be  
20 readily performed by those skilled in the art (see for example, Newton et al, Develop. Dynamics 197: 1-13, 1993).

A binding agent may be a ribosome, with or without a peptide component, an RNA molecule, or a polypeptide. A binding agent may be a polypeptide that comprises a kallikrein 11 or PSA polypeptide sequence, a peptide variant thereof, or a non-peptide mimetic of such a sequence. By way of example a  
25 kallikrein 11 or PSA sequence may be a peptide portion of a kallikrein 11 or PSA that is capable of modulating a function mediated by the kallikrein 11 or PSA.

Antibodies for use in the present invention include but are not limited to monoclonal or polyclonal antibodies, immunologically active fragments (e.g. a Fab or (Fab)<sub>2</sub> fragments), antibody heavy chains, humanized antibodies, antibody light chains, genetically engineered single chain F<sub>v</sub> molecules (Ladner et al,  
30 U.S. Pat. No. 4,946,778), chimeric antibodies, for example, antibodies which contain the binding specificity of murine antibodies, but in which the remaining portions are of human origin, or derivatives, such as enzyme conjugates or labeled derivatives.

Antibodies including monoclonal and polyclonal antibodies, fragments and chimeras, may be prepared using methods known to those skilled in the art. Isolated native or recombinant kallikrein 11 and  
35 PSA may be utilized to prepare antibodies. See, for example, Kohler et al. (1975) Nature 256:495-497; Kozbor et al. (1985) J. Immunol Methods 81:31-42; Cote et al. (1983) Proc Natl Acad Sci 80:2026-2030; and Cole et al. (1984) Mol Cell Biol 62:109-120 for the preparation of monoclonal antibodies; Huse et al. (1989) Science. 246:1275-1281 for the preparation of monoclonal Fab fragments; and, Pound (1998) Immunochemical Protocols, Humana Press, Totowa, N.J for the preparation of phagemid or B-lymphocyte

- 11 -

immunoglobulin libraries to identify antibodies. Antibodies specific for kallikrein 11 and PSA may also be obtained from scientific or commercial sources.

In an embodiment of the invention, antibodies are reactive against kallikrein 11 and PSA if they bind with a  $K_a$  of greater than or equal to  $10^{-7}$  M.

5 **Methods for Detecting Kallikrein 11 and PSA**

A variety of methods can be employed for the diagnostic and prognostic evaluation of prostate cancer involving kallikrein 11 and PSA, and the identification of subjects with a predisposition to such disorders. Such methods may, for example, utilize binding agents (e.g. antibodies) directed against kallikrein 11 and PSA, including peptide fragments. In particular, antibodies may be used, for example, for the  
10 detection of either an over- or an under-abundance of kallikrein 11 and PSA relative to a non-disorder state or the presence of a modified (e.g., less than full length) kallikrein 11 and PSA which correlates with a disorder state, or a progression toward a disorder state.

The invention also contemplates a method for detecting prostate cancer comprising producing a profile of levels of kallikrein 11 and PSA and other markers associated with prostate cancer in cells from a  
15 patient, and comparing the profile with a reference to identify a profile for the test cells indicative of disease.

The methods described herein may be used to evaluate the probability of the presence of malignant or pre-malignant cells, for example, in a group of cells freshly removed from a host. Such methods can be used to detect tumors, quantitate their growth, and help in the diagnosis and prognosis of disease. For example, lower levels of kallikrein 11 and PSA may be indicative of advanced disease, e.g. advanced  
20 prostate cancer. The methods can be used to detect the presence of cancer metastasis, as well as confirm the absence or removal of all tumor tissue following surgery, cancer chemotherapy, and/or radiation therapy. They can further be used to monitor cancer chemotherapy and tumor reappearance.

The methods described herein can be adapted for diagnosing and monitoring prostate cancer by detecting kallikrein 11 and PSA in biological samples from a subject. These applications require that the  
25 amount of kallikrein 11 and PSA quantitated in a sample from a subject being tested be compared to levels quantitated for another sample or an earlier sample from the subject, or levels quantitated for a control sample. Levels for control samples from healthy subjects or prostate cancer subjects may be established by prospective and/or retrospective statistical studies. Healthy subjects who have no clinically evident disease or abnormalities may be selected for statistical studies. Diagnosis may be made by a finding of statistically  
30 different levels of kallikrein 11 and PSA compared to a control sample or previous levels quantitated for the same subject.

Binding agents specific for kallikrein 11 and PSA may be used for a variety of diagnostic and assay applications. There are a variety of assay formats known to the skilled artisan for using a binding agent to detect a target molecule in a sample. (For example, see Harlow and Lane, Antibodies: A Laboratory Manual,  
35 Cold Spring Harbor Laboratory, 1988). In general, the presence or absence of prostate cancer in a subject may be determined by (a) contacting a sample from the subject with binding agents for kallikrein 11 and PSA; (b) detecting in the sample levels of kallikrein 11 and PSA that bind to the binding agents; and (c) comparing the levels of kallikrein 11 and PSA with predetermined standards or cut-off values.

In particular embodiments, the binding agent is an antibody.

- 12 -

In an aspect, the invention provides a diagnostic method for monitoring or diagnosing prostate cancer in a subject by quantitating kallikrein 11 and PSA in a biological sample from the subject comprising reacting the sample with antibodies specific for kallikrein 11 and PSA, which are directly or indirectly labeled with detectable substances and detecting the detectable substances.

5 In an aspect of the invention, a method for detecting prostate cancer is provided comprising:

- (a) obtaining a sample suspected of containing kallikrein 11 and PSA associated with prostate cancer;
- (b) contacting said sample with antibodies that specifically bind kallikrein 11 and PSA under conditions effective to bind the antibodies and form complexes;
- 10 (c) measuring the amount of kallikrein 11 and PSA present in the sample by quantitating the amount of the complexes; and
- (d) comparing the amount of kallikrein 11 and PSA present in the samples with the amount of kallikrein 11 and PSA in a control, wherein a change or significant difference in the amount of kallikrein 11 and PSA in the sample compared with the amount in the control is
- 15 indicative of prostate cancer.

In an embodiment, the invention contemplates a method for monitoring the progression of prostate cancer in an individual, comprising:

- (a) contacting antibodies which bind to kallikrein 11 and PSA with a sample from the individual so as to form complexes comprising the antibodies and kallikrein 11 or PSA in
- 20 the sample;
- (b) determining or detecting the presence or amount of complex formation in the sample;
- (c) repeating steps (a) and (b) at a point later in time; and
- (d) comparing the result of step (b) with the result of step (c), wherein a difference in the amount of complex formation is indicative of disease, disease stage, and/or progression of
- 25 the cancer in said individual.

The amount of complexes may also be compared to a value representative of the amount of the complexes from an individual not at risk of, or afflicted with, prostate cancer at different stages. An decrease in complex formation may be indicative of advanced disease e.g. advanced prostate cancer, or an unfavourable prognosis.

30 Antibodies specifically reactive with kallikrein 11 and PSA, or derivatives, such as enzyme conjugates or labeled derivatives, may be used to detect kallikrein 11 and PSA proteins in various samples (e.g. biological materials). They may be used as diagnostic or prognostic reagents and they may be used to detect abnormalities in the level of kallikrein 11 and PSA expression, or abnormalities in the structure, and/or temporal, tissue, cellular, or subcellular location of kallikrein 11 and PSA. Antibodies may also be

35 used to screen potentially therapeutic compounds *in vitro* to determine their effects on disorders (e.g. prostate cancer) involving a kallikrein 11 and PSA protein, and other conditions. *In vitro* immunoassays may also be used to assess or monitor the efficacy of particular therapies.

Antibodies may be used in any known immunoassays that rely on the binding interaction between antigenic determinants of kallikrein 11 and PSA protein and the antibodies. Immunoassay procedures for *in*

- 13 -

*vitro* detection of antigens in fluid samples are also well known in the art. [See for example, Paterson et al., Int. J. Can. 37:659 (1986) and Burchell et al., Int. J. Can. 34:763 (1984) for a general description of immunoassay procedures]. Qualitative and/or quantitative determinations of kallikrein 11 and PSA in a sample may be accomplished by competitive or non-competitive immunoassay procedures in either a direct or indirect format. Detection of kallikrein 11 and PSA using antibodies can be done utilizing immunoassays which are run in either the forward, reverse or simultaneous modes. Examples of immunoassays are radioimmunoassays (RIA), enzyme immunoassays (e.g. ELISA), immunofluorescence, immunoprecipitation, latex agglutination, hemagglutination, histochemical tests, and sandwich (immunometric) assays. These terms are well understood by those skilled in the art. A person skilled in the art will know, or can readily discern, other immunoassay formats without undue experimentation. The antibodies may be used to detect and quantify kallikrein 11 and PSA in a sample in order to diagnose and treat pathological states.

In particular, the antibodies may be used in immunohistochemical analyses, for example, at the cellular and sub-subcellular level, to detect kallikrein 11 and PSA proteins, to localize them to particular prostate tumor cells and tissues, and to specific subcellular locations, and to quantitate the level of expression.

Immunohistochemical methods for the detection of antigens in tissue samples are well known in the art. For example, immunohistochemical methods are described in Taylor, Arch. Pathol. Lab. Med. 102:112 (1978). Briefly, in the context of the present invention, a tissue sample obtained from a subject suspected of having a prostate-related problem is contacted with antibodies, preferably monoclonal antibodies recognizing kallikrein 11. The site at which the antibodies are bound is determined by selective staining of the sample by standard immunohistochemical procedures. The same procedure may be repeated on the same sample using other antibodies that recognize kallikrein 11. Alternatively, a sample may be contacted with antibodies against kallikrein 11 and antibodies against PSA simultaneously, provided that the antibodies are labeled differently or are able to bind to a different label. In one embodiment of the present invention, the tissue sample is obtained from the prostate of a patient. The prostate tissue sample may be a normal prostate tissue, a cancer prostate tissue or a benign prostatic hyperplasia tissue.

In a sandwich immunoassay of the invention mouse polyclonal/monoclonal antibodies specific for kallikrein 11 and PSA and rabbit polyclonal/monoclonal antibodies specific for kallikrein 11 and PSA are utilized.

Antibodies specific for kallikrein 11 and PSA may be labelled with a detectable substance and localised in biological samples based upon the presence of the detectable substance. Examples of detectable substances include, but are not limited to, the following: radioisotopes (e.g.,  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{35}\text{S}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ), fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), luminescent labels such as luminol; enzymatic labels (e.g., horseradish peroxidase, beta-galactosidase, luciferase, alkaline phosphatase, acetylcholinesterase), biotinyl groups (which can be detected by marked avidin e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or calorimetric methods); predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, labels

- 14 -

are attached via spacer arms of various lengths to reduce potential steric hindrance. Antibodies may also be coupled to electron dense substances, such as ferritin or colloidal gold, which are readily visualised by electron microscopy.

One of the ways an antibody can be detectably labeled is to link it directly to an enzyme. The enzyme when later exposed to its substrate will produce a product that can be detected. Examples of detectable substances that are enzymes are horseradish peroxidase, beta-galactosidase, luciferase, alkaline phosphatase, acetylcholinesterase, malate dehydrogenase, ribonuclease, urease, catalase, glucose-6-phosphate, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate, triose phosphate isomerase, asparaginase, glucose oxidase, and acetylcholine esterase.

For increased sensitivity in an immunoassay system a fluorescence-emitting metal atom such as Eu (europium) and other lanthanides can be used. These can be attached to the desired molecule by means of metal-chelating groups such as DTPA or EDTA.

A bioluminescent compound may also be used as a detectable substance. Bioluminescence is a type of chemiluminescence found in biological systems where a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent molecule is determined by detecting the presence of luminescence. Examples of bioluminescent detectable substances are luciferin, luciferase and aequorin.

Indirect methods may also be employed in which the primary antigen-antibody reaction is amplified by the introduction of a second antibody, having specificity for the antibody reactive against kallikrein 11 and PSA. By way of example, if the antibody having specificity against kallikrein 11 and PSA is a rabbit IgG antibody, the second antibody may be goat anti-rabbit IgG, Fc fragment specific antibody labelled with a detectable substance as described herein.

Methods for conjugating or labelling the antibodies discussed above may be readily accomplished by one of ordinary skill in the art. (See for example Inman, *Methods In Enzymology*, Vol. 34, *Affinity Techniques*, Enzyme Purification: Part B, Jakoby and Wichek (eds.), Academic Press, New York, p. 30, 1974; and Wilchek and Bayer, "The Avidin-Biotin Complex in Bioanalytical Applications," *Anal. Biochem.* 171:1-32, 1988 re methods for conjugating or labelling the antibodies with enzyme or ligand binding partner).

Cytochemical techniques known in the art for localizing antigens using light and electron microscopy may be used to detect a kallikrein 11 and PSA protein. Generally, an antibody may be labeled with a detectable substance and a kallikrein 11 and PSA protein may be localised in tissues and cells based upon the presence of the detectable substance.

In the context of the methods of the invention, the sample, binding agents (e.g. antibodies) or kallikrein 11 and PSA may be immobilized on a carrier or support. Examples of suitable carriers or supports are agarose, cellulose, nitrocellulose, dextran, Sephadex, Sepharose, liposomes, carboxymethyl cellulose, polyacrylamides, polystyrene, gabbros, filter paper, magnetite, ion-exchange resin, plastic film, plastic tube, glass, polyamine-methyl vinyl-ether-maleic acid copolymer, amino acid copolymer, ethylene-maleic acid copolymer, nylon, silk, etc. The support material may have any possible configuration including spherical (e.g. bead), cylindrical (e.g. inside surface of a test tube or well, or the external surface of a rod), or flat (e.g.



- 15 -

sheet, test strip). Thus, the carrier may be in the shape of, for example, a tube, test plate, well, beads, disc, sphere, etc. The immobilized material may be prepared by reacting the material with a suitable insoluble carrier using known chemical or physical methods, for example, cyanogen bromide coupling. Binding agents (e.g. antibodies) may be indirectly immobilized using second binding agents specific for the first binding agent. For example, mouse antibodies specific for kallikrein 11 and PSA may be immobilized using sheep anti-mouse IgG Fc fragment specific antibody coated on the carrier or support.

Where a radioactive label is used as a detectable substance, a kallikrein 11 and PSA protein may be localized by radioautography. The results of radioautography may be quantitated by determining the density of particles in the radioautographs by various optical methods, or by counting the grains.

Time-resolved fluorometry may be used to detect a fluorescent signal, label, or detectable substance. For example, the method described in Christopoulos TK and Diamandis EP Anal. Chem., 1992:64:342-346 may be used with a conventional time-resolved fluorometer.

According to an embodiment of the invention, an immunoassay for detecting kallikrein 11 in a biological sample comprises contacting an amount of a first agent that specifically binds to kallikrein 11 in the sample under a condition that allows the formation of a first complex comprising the first agent and kallikrein 11 and determining the presence or amount of the complex as a measure of the amount of the kallikrein 11 contained in the sample. In another embodiment, an immunoassay is provided for detecting kallikrein 11 and total PSA in a sample comprising the additional step of contacting an amount of a second agent which specifically binds to PSA with the sample under a condition that allows the formation of a second complex comprising the second agent and PSA, and determining the presence or amount of the second complex as a measure of the amount of the PSA contained in the sample. In an alternate method, a sample may be contacted by the first and the second agents simultaneously, provided that the agents are labeled differently or are capable of binding to different labels.

In accordance with an embodiment of the invention, a method is provided wherein kallikrein 11 and PSA antibodies are directly or indirectly labelled with enzymes, substrates for the enzymes are added wherein the substrates are selected so that the substrates, or a reaction product of an enzyme and substrate, form fluorescent complexes with a lanthanide metal (e.g. europium, terbium, samarium, and dysprosium, preferably europium and terbium). A lanthanide metal is added and kallikrein 11 and PSA are quantitated in the sample by measuring fluorescence of the fluorescent complexes. Enzymes are selected based on the ability of a substrate of the enzyme, or a reaction product of the enzyme and substrate, to complex with lanthanide metals such as europium and terbium. Suitable enzymes and substrates that provide fluorescent complexes are described in U.S. Patent No. 5,312,922 to Diamandis. Examples of suitable enzymes include alkaline phosphatase and  $\beta$ -galactosidase. Preferably, the enzyme is alkaline phosphatase.

Examples of enzymes and substrates for enzymes that provide such fluorescent complexes are described in U.S. Patent No. 5,312,922 to Diamandis. By way of example, when the antibody is directly or indirectly labelled with alkaline phosphatase the substrate employed in the method may be 4-methylumbelliferyl phosphate, 5-fluorosalicyl phosphate, or diflunisal phosphate. The fluorescence intensity of the complexes is typically measured using a time-resolved fluorometer e.g. a CyberFluor 615 Imunoanalyzer (Nordion International, Kanata, Ontario).

- 16 -

Antibodies specific for kallikrein 11 and PSA may also be indirectly labelled with enzymes. For example, an antibody may be conjugated to one partner of a ligand binding pair, and the enzyme may be coupled to the other partner of the ligand binding pair. Representative examples include avidin-biotin, and riboflavin-riboflavin binding protein. In an embodiment, an antibody specific for the anti-kallikrein 11 and PSA antibody is labelled with an enzyme.

In accordance with an embodiment, the present invention provides means for determining kallikrein 11 and PSA in a serum sample by measuring kallikrein 11 and PSA by immunoassay. It will be evident to a skilled artisan that a variety of immunoassay methods can be used to measure kallikrein 11 and PSA in serum. In general, a kallikrein 11 and PSA immunoassay method may be competitive or noncompetitive. Competitive methods typically employ an immobilized or immobilizable antibody to kallikrein 11 and PSA (anti-kallikrein 11 and anti-PSA) and a labeled form of kallikrein 11 and PSA. Sample kallikrein 11 and PSA and labeled kallikrein 11 and PSA compete for binding to anti-kallikrein 11 and PSA. After separation of the resulting labeled kallikrein 11 and PSA that has become bound to anti-kallikrein 11 and PSA (bound fraction) from that which has remained unbound (unbound fraction), the amount of the label in either bound or unbound fraction is measured and may be correlated with the amount of kallikrein 11 and PSA in the test sample in any conventional manner, e.g., by comparison to a standard curve.

In an aspect, a non-competitive method is used for the determination of kallikrein 11 and PSA, with the most common method being the "sandwich" method. In this assay, two anti- kallikrein 11 and two PSA antibodies are employed. One of the anti- kallikrein 11 and one of the PSA antibodies is directly or indirectly labeled (sometimes referred to as the "detection antibody") and the other is immobilized or immobilizable (sometimes referred to as the "capture antibody"). The capture and detection antibodies can be contacted simultaneously or sequentially with the test sample. Sequential methods can be accomplished by incubating the capture antibody with the sample, and adding the detection antibody at a predetermined time thereafter (sometimes referred to as the "forward" method); or the detection antibody can be incubated with the sample first and then the capture antibody added (sometimes referred to as the "reverse" method). After the necessary incubation(s) have occurred, to complete the assay, the capture antibody may be separated from the liquid test mixture, and the label may be measured in at least a portion of the separated capture antibody phase or the remainder of the liquid test mixture. Generally it is measured in the capture antibody phase since it comprises kallikrein 11 and PSA bound by ("sandwiched" between) the capture and detection antibodies. In another embodiment, the label may be measured without separating the capture antibody and liquid test mixture.

In a typical two-site immunometric assay for kallikrein 11 and PSA, one or both of the capture and detection antibodies are polyclonal antibodies or one or both of the capture and detection antibodies are monoclonal antibodies (i.e. polyclonal/polyclonal, monoclonal/monoclonal, or monoclonal/polyclonal). The label used in the detection antibody can be selected from any of those known conventionally in the art. The label may be an enzyme or a chemiluminescent moiety, but it can also be a radioactive isotope, a fluorophor, a detectable ligand (e.g., detectable by a secondary binding by a labeled binding partner for the ligand), and the like. Preferably the antibody is labelled with an enzyme which is detected by adding a substrate that is selected so that a reaction product of the enzyme and substrate forms fluorescent complexes. The capture

- 17 -

antibody may be selected so that it provides a means for being separated from the remainder of the test mixture. Accordingly, the capture antibody can be introduced to the assay in an already immobilized or insoluble form, or can be in an immobilizable form, that is, a form which enables immobilization to be accomplished subsequent to introduction of the capture antibody to the assay. An immobilized capture antibody may comprise an antibody covalently or noncovalently attached to a solid phase such as a magnetic particle, a latex particle, a microtiter plate well, a bead, a cuvette, or other reaction vessel. An example of an immobilizable capture antibody is antibody which has been chemically modified with a ligand moiety, e.g., a hapten, biotin, or the like, and which can be subsequently immobilized by contact with an immobilized form of a binding partner for the ligand, e.g., an antibody, avidin, or the like. In an embodiment, the capture antibody may be immobilized using a species specific antibody for the capture antibody that is bound to the solid phase.

A particular sandwich immunoassay method of the invention employs antibodies reactive against kallikrein 11 and PSA, antibodies having specificity against antibodies reactive against kallikrein 11 and PSA labelled with an enzymatic label, and a fluorogenic substrate for the enzyme. In an embodiment, the enzyme is alkaline phosphatase (ALP) and the substrate is 5-fluorosalicyl phosphate. ALP cleaves phosphate out of the fluorogenic substrate, 5-fluorosalicyl phosphate, to produce 5-fluorosalicylic acid (FSA). 5-Fluorosalicylic acid can then form a highly fluorescent ternary complex of the form FSA-Tb<sup>3+</sup>-EDTA, which can be quantified by measuring the Tb<sup>3+</sup> fluorescence in a time-resolved mode. Fluorescence intensity is measured using a time-resolved fluorometer as described herein.

The above-described immunoassay methods and formats are intended to be exemplary and are not limiting.

The invention also contemplates the methods described herein using multiple markers for prostate cancer. Therefore, the invention contemplates a method for analyzing a biological sample for the presence of kallikrein 11 and PSA and other markers that are specific indicators of prostate cancer. The methods described herein may be modified by including reagents to detect the markers, or nucleic acids for the markers. The methods described herein may also include reagents to detect KLK11. Techniques for detecting nucleic acid such as polymerase chain reaction (PCR) and hybridization assays are well known in the art.

The use of kallikrein 11 alone, and kallikrein 11 in combination with PSA may lead to improved discrimination between BPH and prostate cancer. The mathematical combination of the amount of kallikrein 11 and PSA may be used as a serum marker or as an immunohistological marker to help distinguish prostate cancer from BPH. Alternatively, kallikrein 11 alone may be used as a serum marker or as an immunohistological marker for distinguishing BPH from prostate cancer. The term "mathematical combination" as used herein refers to any mathematical calculation of the amount of kallikrein 11 and PSA. In an aspect of the invention, the mathematical combination is a ratio. The ratio of kallikrein 11 and PSA in a sample may be determined by comparing the amount of kallikrein 11 to the amount of total PSA in the sample. Thus, the mathematical combination may be the ratio of kallikrein 11: total PSA or the inverse thereof.

The mathematical combination may be compared to the mathematical combination of a standard in order to determine the presence of BPH or prostate cancer in the subject. In a particular embodiment of the

- 18 -

invention a kallikrein 11: total PSA ratio greater than 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.10, 0.11, or 0.12 is potentially indicative of BPH while a kallikrein 11: total PSA ratio less than 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.10, 0.11, or 0.12 is potentially indicative of prostate cancer. In a more particular embodiment, a kallikrein 11: total PSA ratio greater than 0.05 is potentially indicative of BPH while a  
 5 kallikrein 11: total PSA ratio less than 0.05 is potentially indicative of prostate cancer.

Accordingly, another aspect of the present invention provides a method for determining the presence of BPH or prostate cancer in a subject comprising the steps of:

- (a) providing a first binding agent that specifically binds to kallikrein 11;
- (b) providing a second binding agent that specifically binds to PSA;
- 10 (c) contacting the first agent and the second agent with a sample from the subject under a condition that allows the formation of a first complex comprising the first agent and kallikrein 11 and a second complex comprising the second agent and the PSA;
- (d) detecting or determining the presence or amount of the first and second complexes;
- (e) mathematically combining the amount of the first and the second complexes or the amount  
 15 of kallikrein 11; and
- (f) relating the combination to the presence of BPH or prostate cancer in the subject.

In accordance with embodiments of the present invention, the agents comprise antibodies. Preferably, the first and the second agents are directly or indirectly labeled with different detectable substances to form respective complexes that may be detected separately. Alternatively, the sample may be  
 20 contacted with one agent first so that either kallikrein 11 may be detected first, then the same sample may be contacted with another agent in order to detect PSA.

In one embodiment of the present invention, the sample used in the diagnostic method may be a sample of human physiological fluid such as, but not limited to, serum, seminal plasma, urine and plasma, preferably serum. In another embodiment, the sample may be tissue specimen from the prostate of a patient.

25 In a particular embodiment, the combination is a ratio of the first complex to the second complex, or the inverse thereof.

In particular methods of the invention, patients with BPH or prostate cancer can be identified in patients with total PSA between about 4-10ng/ml.

In particular methods of the invention, patients with BPH or prostate cancer can be identified in  
 30 patients with total PSA less than about 4 ng/ml.

The methods of the invention may utilize commercially available kits for detecting or quantifying PSA in a sample. For example the Immulite PSA (Diagnostic Products Corporation, Cal, USA) assay may be used to measure PSA in a sample.

35 Since kallikrein 11 alone may be used as a serum marker or as an immunohistological marker for distinguishing BPH from prostate cancer, the present invention also provides a diagnostic method for distinguishing BPH from prostate cancer by detecting and determining the amount of kallikrein 11 in a sample. The amount of kallikrein 11 may be determined in tissue samples by immunohistochemical methods and/or in patient fluid samples by *in vitro* immunoassay procedures described herein.

### Computer Systems

Computer readable media comprising kallikrein 11 and PSA and optionally other markers of prostate cancer is also provided. "Computer readable media" refers to any medium that can be read and accessed directly by a computer, including but not limited to magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. Thus, the invention contemplates computer readable medium having recorded thereon markers identified for patients and controls.

"Recorded" refers to a process for storing information on computer readable medium. The skilled artisan can readily adopt any of the presently known methods for recording information on computer readable medium to generate manufactures comprising information on kallikrein 11 and PSA and optionally other prostate cancer markers.

A variety of data processor programs and formats can be used to store information on kallikrein 11 and PSA and other prostate cancer markers on computer readable medium. For example, the information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and MicroSoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. Any number of dataprocessor structuring formats (e.g., text file or database) may be adapted in order to obtain computer readable medium having recorded thereon the marker information.

By providing the marker information in computer readable form, one can routinely access the information for a variety of purposes. For example, one skilled in the art can use the information in computer readable form to compare marker information obtained during or following therapy with the information stored within the data storage means.

The invention provides a medium for holding instructions for performing a method for determining whether a patient has prostate cancer or a pre-disposition to prostate cancer, comprising determining the presence, absence, or amount of kallikrein 11 and PSA and optionally other prostate cancer markers, and based on the presence, absence or amount of kallikrein 11 and PSA and optionally other markers, determining whether the patient has prostate cancer, BPH, or a pre-disposition to prostate cancer, and optionally recommending treatment for the condition.

The invention also provides in an electronic system and/or in a network, a method for determining whether a subject has prostate cancer or a pre-disposition to prostate cancer, comprising determining the presence, absence, or amount of kallikrein 11 and PSA and optionally other prostate cancer markers, and based on the presence, absence, or amount of the kallikrein 11 and PSA and optionally other markers, determining whether the subject has prostate cancer, BPH, or a pre-disposition to prostate cancer, and optionally recommending treatment for the condition.

The invention further provides in a network, a method for determining whether a subject has prostate cancer or a pre-disposition to prostate cancer comprising: (a) receiving phenotypic information on the subject and information on kallikrein 11 and PSA and optionally other prostate cancer markers associated with samples from the subject; (b) acquiring information from the network corresponding to the kallikrein

- 20 -

11 and PSA and optionally other markers; and (c) based on the phenotypic information and information on the kallikrein 11 and PSA and optionally other markers determining whether the subject has prostate cancer, BPH or a pre-disposition to prostate cancer; and (d) optionally recommending treatment for the condition.

5 The invention still further provides a system for identifying selected records that identify a prostate cancer cell or tissue. A system of the invention generally comprises a digital computer; a database server coupled to the computer; a database coupled to the database server having data stored therein, the data comprising records of data comprising kallikrein 11 and PSA and optionally other prostate cancer markers, or nucleic acids encoding same, and a code mechanism for applying queries based upon a desired selection  
10 criteria to the data file in the database to produce reports of records which match the desired selection criteria.

In an aspect of the invention a method is provided for detecting prostate cancer tissue or cells using a computer having a processor, memory, display, and input/output devices, the method comprising the steps of:

- 15 (a) creating records of kallikrein 11 and PSA including mathematical combinations thereof, and optionally other prostate cancer markers isolated from a sample suspected of containing prostate cancer cells or tissue;
- (b) providing a database comprising records of data comprising kallikrein 11 and PSA including mathematical combinations thereof, and optionally other prostate cancer markers; and
- 20 (c) using a code mechanism for applying queries based upon a desired selection criteria to the data file in the database to produce reports of records of step (a) which provide a match of the desired selection criteria of the database of step (b) the presence of a match being a positive indication that the markers of step (a) have been isolated from cells or tissue that are prostate cancer cells or tissue.

25 The invention contemplates a business method for determining whether a subject has prostate cancer or a pre-disposition to prostate cancer comprising: (a) receiving phenotypic information on the subject and information on kallikrein 11 and PSA and optionally other prostate cancer markers associated with samples from the subject; (b) acquiring information from a network corresponding to kallikrein 11 and PSA and optionally other markers; and (c) based on the phenotypic information, information on kallikrein 11 and  
30 PSA and optionally other markers, and acquired information, determining whether the subject has prostate cancer or a pre-disposition to prostate cancer; and (d) optionally recommending treatment for the prostate cancer or pre-condition.

In an aspect of the invention, the computer systems, components, and methods described herein are used to monitor disease or determine the stage of disease.

#### 35 Screening Methods

The invention also contemplates methods for evaluating test agents or compounds for their ability to inhibit prostate cancer or potentially contribute to prostate cancer. Test agents and compounds include but are not limited to peptides such as soluble peptides including Ig-tailed fusion peptides, members of random peptide libraries and combinatorial chemistry-derived molecular libraries made of D- and/or L-configuration

- 21 -

amino acids, phosphopeptides (including members of random or partially degenerate, directed phosphopeptide libraries), antibodies [e.g. polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, single chain antibodies, fragments, (e.g. Fab, F(ab)<sub>2</sub>, and Fab expression library fragments, and epitope-binding fragments thereof]], and small organic or inorganic molecules. The agents or compounds may be endogenous physiological compounds or natural or synthetic compounds.

The invention provides a method for assessing the potential efficacy of a test agent for inhibiting prostate cancer in a patient, the method comprising comparing:

- (a) levels of kallikrein 11 and PSA including mathematical combinations thereof, and optionally other prostate cancer markers in a first sample obtained from a patient and exposed to the test agent; and
- (b) levels of kallikrein 11 and PSA including mathematical combinations thereof, and optionally other markers in a second sample obtained from the patient, wherein the sample is not exposed to the test agent, wherein a significant difference in the levels of expression of kallikrein 11 and PSA including mathematical combinations thereof, and optionally the other markers in the first sample, relative to the second sample, is an indication that the test agent is potentially efficacious for inhibiting prostate cancer in the patient.

The first and second samples may be portions of a single sample obtained from a patient or portions of pooled samples obtained from a patient.

In an embodiment, the levels of expression of kallikrein 11 and PSA including mathematical combinations thereof, in the first sample are significantly higher relative to the second sample.

In an aspect, the invention provides a method of selecting an agent for inhibiting prostate cancer in a patient comprising:

- (a) obtaining a sample from the patient;
- (b) separately maintaining aliquots of the sample in the presence of a plurality of test agents;
- (c) comparing kallikrein 11 and PSA including mathematical combinations thereof and optionally other prostate cancer markers, in each of the aliquots; and
- (d) selecting one of the test agents which alters the levels of kallikrein 11 and PSA including mathematical combinations thereof and optionally other prostate cancer markers in the aliquot containing that test agent, relative to other test agents.

In an embodiment, the levels of kallikrein 11 and PSA including mathematical combinations thereof are significantly higher in the presence of the selected test agent.

Still another aspect of the present invention provides a method of conducting a drug discovery business comprising:

- (a) providing one or more methods or assay systems for identifying agents that inhibit prostate cancer in a patient;
- (b) conducting therapeutic profiling of agents identified in step (a), or further analogs thereof, for efficacy and toxicity in animals; and
- (c) formulating a pharmaceutical preparation including one or more agents identified in step (b) as having an acceptable therapeutic profile.

- 22 -

In certain embodiments, the subject method can also include a step of establishing a distribution system for distributing the pharmaceutical preparation for sale, and may optionally include establishing a sales group for marketing the pharmaceutical preparation.

The invention also contemplates a method of assessing the potential of a test compound to contribute to prostate cancer comprising:

- (a) maintaining separate aliquots of cells or tissues from a patient with prostate cancer in the presence and absence of the test compound; and
- (b) comparing kallikrein 11 and PSA including mathematical combinations thereof and optionally other prostate cancer markers in each of the aliquots.

A significant difference between the levels of the markers in the aliquot maintained in the presence of (or exposed to) the test compound relative to the aliquot maintained in the absence of the test compound, indicates that the test compound possesses the potential to contribute to prostate cancer. In an embodiment, the levels of kallikrein 11 and PSA including mathematical combinations thereof are lower in the presence of the test compound.

In an aspect the invention provides a method of inhibiting prostate cancer in a patient, the method comprising (a) obtaining a sample comprising cells affected by prostate cancer from the patient; (b) separately maintaining aliquots of the sample in the presence of a plurality of test agents; (c) comparing levels of kallikrein 11 and PSA or a mathematical combination thereof in each of the aliquots; and (d) administering to the patient at least one of the test agents which alters kallikrein 11 and PSA or a mathematical combination thereof in the aliquot containing that test agent, relative to other test agents.

#### Kits

The invention contemplates kits for carrying out the methods of the invention. Such kits typically comprise two or more components required for performing a diagnostic assay. Components include but are not limited to compounds, reagents, containers, and/or equipment.

The methods described herein may be performed by utilizing pre-packaged diagnostic kits comprising at least binding agents (e.g. antibodies) described herein, which may be conveniently used, e.g. in clinical settings, to screen and diagnose patients, and to screen and identify those individuals afflicted with prostate cancer or BPH or exhibiting a predisposition to such condition.

In an embodiment, a container with a kit comprises binding agents as described herein. By way of example, the kit may contain antibodies specific for kallikrein 11 and PSA and optionally other prostate cancer markers, antibodies against the antibodies labelled with enzymes, and substrates for the enzymes. The kit may also contain microtiter plate wells, standards, assay diluent, wash buffer, adhesive plate covers, and/or instructions for carrying out a method of the invention using the kit.

In an aspect of the invention, the kit includes antibodies or antibody fragments which bind specifically to epitopes of kallikrein 11 and PSA and optionally other prostate cancer markers, and means for detecting binding of the antibodies to epitopes associated with prostate cancer, either as concentrates (including lyophilized compositions), which may be further diluted prior to use or at the concentration of use, where the vials may include one or more dosages. Where the kits are intended for *in vivo* use, single dosages may be provided in sterilized containers, having the desired amount and concentration of agents.



- 23 -

Containers that provide a formulation for direct use, usually do not require other reagents.

In particular, the invention provides a kit for determining the presence of BPH or prostate comprising a known amount of a first binding agent that specifically binds to kallikrein 11 wherein the first binding agent comprises a detectable substance, or it binds directly or indirectly to a detectable substance.

5 Another aspect of the present invention also provides a kit for determining the presence of BPH or prostate cancer in a sample. The kit includes:

- (a) a known amount of a first agent which specifically binds to kallikrein 11, and
- (b) a known amount of a second agent which specifically binds to a PSA,

10 wherein the first and the second agents, respectively, comprise a detectable label or bind to a detectable label.

The agents may be antibodies, particularly monoclonal antibodies. Preferably, the first and the second agents, respectively, comprise a different detectable substance or bind to a different detectable substance.

15 The reagents suitable for applying the screening methods of the invention to evaluate compounds may be packaged into convenient kits described herein providing the necessary materials packaged into suitable containers.

Thus, the invention relates to a kit for assessing the suitability of each of a plurality of test compounds for inhibiting prostate cancer in a patient. The kit comprises reagents for assessing kallikrein 11 and PSA and optionally a plurality of test agents or compounds.

20 The invention contemplates a kit for assessing the presence of prostate cancer cells, wherein the kit comprises antibodies specific for kallikrein 11 and PSA, and optionally antibodies specific for other markers associated with prostate cancer.

Additionally the invention provides a kit for assessing the potential of a test compound to contribute to prostate cancer. The kit comprises prostate cancer cells and reagents for assessing kallikrein 11 and PSA, 25 and optionally other markers associated with endocrine cancer.

The following examples are intended to illustrate, but not to limit, the scope of the invention. While such examples are typical of those that might be used, other procedures known to those skilled in the art may alternatively be utilized. Indeed, those of ordinary skill in the art can readily envision and produce further embodiments, based on the teachings herein, without undue experimentation.

30 **Example 1**

hK11, total PSA and % free PSA, was analyzed in a total of 150 serum samples from men with histologically confirmed benign prostatic hyperplasia (BPH; N = 64) or prostatic cancer (CaP; N = 86). Total and free PSA levels were measured by the Immulite PSA assay and hK11 levels were measured by an immunofluorometric assay. Serum hK11 levels and the hK11/total PSA ratio were both significantly lower 35 in CaP patients than in BPH. In the subgroup of patients with % free PSA less than 20, an additional 54 % of BPH patients could have avoided biopsies using the hK11/total PSA ratio. ROC curve analysis demonstrated that the hK11/ total PSA ratio (AUC = 0.81) and % free PSA (AUC = 0.82) were much stronger predictors of prostate cancer than total PSA (AUC = 0.69). Thus, the hK11/total PSA ratio is a useful tumor marker for prostate cancer and may be combined with % free PSA to further significantly reduce the number of

unnecessary prostatic biopsies.

### **Materials and Methods**

#### **Study population**

5 Included in this study were serum samples from 150 male patients, 64 with benign prostate hyperplasia (BPH) (median age 65) and 86 with prostate cancer (CaP) (median age 62), all histologically confirmed by biopsy. All patients were from the Department of Urology at University Hospital Charité, Humboldt University, Berlin, Germany. Total and %free PSA, measured by the Immulite method (Diagnostic Products Corp., San Diego, CA) were available for all samples. All sera were stored at  $-80^{\circ}\text{C}$  until use.

10 The samples were collected with informed consent and the study was approved by the Institutional Review Board of the University Hospital Charité, Humboldt University, Berlin, Germany.

#### **Production of monoclonal and polyclonal antibodies against hK11**

15 Purified recombinant hK11 protein was used to immunize rabbits and mice, to raise polyclonal and monoclonal antibodies as described elsewhere<sup>11</sup>. hK11 was produced by expression in Pichia Pastoris (Invitrogen) and purified by SP Sepharose cation exchange column and reverse phase liquid chromatography.

#### **Development of immunofluorometric assay for hK11**

20 In this study, the hK11 assay described previously was used, with minor modifications<sup>11</sup>. In brief, sheep anti-mouse IgG (Jackson ImmunoResearch) diluted in coating buffer (containing 50mM Tris, pH 7.8) was dispensed into a 96-well white polystyrene microtiter plate (500 ng/ 100  $\mu\text{l}$ /well) and incubated overnight at room temperature. The plates were then washed three times with washing buffer that contains 9 g/L NaCl and 0.5 g/L Tween 20 in 10mM Tris, pH 7.8. One hundred  $\mu\text{l}$  of anti-hK11 monoclonal antibody (100ng) diluted in 6% BSA were added to each well and incubated with orbital shaking for 2 hours at room temperature. The plates were washed 6 times with wash buffer. Fifty  $\mu\text{l}$  of hK11 calibrators (recombinant hK11 in 6% BSA) or samples were applied into each well along with 50  $\mu\text{l}$  of 6% BSA. The plates were 25 incubated for 2 hours on an orbital shaker. After this step, the previously described procedure was used<sup>11</sup>. All samples were analyzed in triplicate.

#### **Statistical analysis**

30 The analysis of differences between measured or calculated parameters, in the two groups, were performed with the non-parametric Mann-Whitney U test. Relationships between different variables were assessed by Spearman correlation coefficient. Receiver operating characteristic (ROC) curves were constructed for total PSA, % free PSA and hK11/total PSA ratio, by plotting sensitivity versus (1-specificity) and the area under the ROC curves (AUC) were analyzed by the Hanley and McNeil method.

#### **Results**

35 hK11 levels were measured in 150 serum samples from 86 patients with prostate cancer and 64 patients with benign prostate hyperplasia with known total and % free PSA values. Descriptive statistics are summarized in Tables 1 and 2.

Total PSA values ranged from 0.17 to 26.2  $\mu\text{g/L}$  in BPH patients, with a mean  $\pm$  SE of  $5.61 \pm 0.63$   $\mu\text{g/L}$  and from 0.35 to 48.0  $\mu\text{g/L}$  in CaP patients, with the mean  $\pm$  SE of  $9.34 \pm 0.82$   $\mu\text{g/L}$ . Percent free PSA

levels ranged from 0.85 to 64.0 (mean  $\pm$  SE =  $22.3 \pm 0.16$ ) and from 1.9 to 47.9 (mean  $\pm$  SE =  $10.9 \pm 0.8$ ) in patients with BPH and CaP, respectively. hK11 concentrations ranged from 0.0 to 2.33  $\mu\text{g/L}$  (mean  $\pm$  SE of  $0.41 \pm 0.05$   $\mu\text{g/L}$ ) in BPH patients and from 0.0 to 0.72  $\mu\text{g/L}$  (mean  $\pm$  SE of  $0.15 \pm 0.02$   $\mu\text{g/L}$ ) in CaP patients. The distributions of hK11 between the two groups were significantly different ( $p < 0.001$ ). The mean value  $\pm$  SE of hK11/total PSA ratio was  $0.14 \pm 0.03$  (range 0.00 – 1.28) in patients with BPH and  $0.028 \pm 0.005$  (range 0.00 – 0.31) in patients with CaP. The distribution of hK11/total PSA ratio was significantly lower in CaP patients than in BPH patients ( $p < 0.001$ ). At 90 % sensitivity (hK11/total PSA ratio of 0.06) the specificity is 48.5% (Figure 1). Furthermore, the hK11/total PSA ratio was analyzed in the subgroup of patients with % free PSA  $< 20$  (these patients would have undergone biopsy based on % free PSA). At a cutoff-point of 0.05 (90% sensitivity), specificity was 54%. Nineteen of these 35 patients could have avoided biopsy based on this criterion (Figure 2).

A weak positive correlation was found between hK11 and total PSA levels in the group of BPH patients (Spearman correlation coefficient  $r_s = 0.29$ ,  $p = 0.017$ ). However, in CaP patients, a significant correlation between hK11 and other variables was not observed.

The discriminatory value of the hK11/total PSA ratio was investigated in relation to the total PSA and % free PSA values, by ROC curve analysis. The hK11/total PSA ratio, overall, had about the same discriminatory potential as % free PSA (AUC = 0.81, 95% CI = 0.74 – 0.88 for the hK11/total PSA ratio and AUC = 0.82, 95% CI = 0.76 – 0.89 for the % free PSA) (Figure 3).

#### Discussion

PSA is widely used as a tumor marker for prostate cancer, but the discrimination between BPH and CaP with this test is not possible due to the lack of specificity. It has been reported that PSA molecular forms, including free PSA, can contribute to better discrimination<sup>4,12-14</sup>. hK2 is also useful for the differential diagnosis for prostate cancer in the PSA range of 2.5 to 10  $\mu\text{g/L}$ <sup>4,15</sup>.

Recently, an immunoassay was developed for hK11 and relatively large amounts of hK11 was found in seminal plasma and prostatic tissues<sup>11</sup>.

In this study, it is demonstrated for the first time that hK11 levels in serum are significantly lower in CaP patients than in BPH patients. The hK11/total PSA ratio is also significantly lower in CaP than in BPH. Using the hK11/total PSA ratio, approximately 50% of biopsies could be avoided that could have not been avoided by the % free PSA test (% free PSA  $< 20$ ). These results demonstrate for the first time that the combination of % free PSA and the hK11/total PSA ratio could contribute to better discrimination between BPH and CaP patients. Out of 64 patients with BPH, and at 90 % sensitivity, 29 patients (45%) would have avoided biopsy by % free PSA testing ( $> 20$  % free PSA). For the remaining 35 patients, another 19 could have avoided biopsy by the hK11/total PSA ratio ( $> 0.05$ ). When the two tests are combined, 48 BPH patients (80 %) would have avoided biopsy at about 90 % sensitivity (Figure 4).

ROC curve analysis demonstrated that the hK11/total PSA ratio had about the same discriminatory value as % free PSA, suggesting that hK11 could be an additional marker for prostate cancer.

#### Conclusions

The potential clinical utility of serum hK11 analysis for the differential diagnosis of prostate cancer

was investigated and it was determined that the hK11/total PSA ratio could be a useful novel tumor marker.

#### Example 2

Human kallikrein 11, total PSA, and %free PSA were analyzed using the procedure described in Example 1 in serum samples from 405 male patients, 247 with benign prostatic hyperplasia (median age 66) and 157 with prostate cancer (median age 64), histologically confirmed by biopsy in all cases. Descriptive statistics are described in Table 3. The logistic regression analysis of BPH and CaP patients for predicting the presence of prostate cancer is shown in Table 4. A comparison of the sensitivity and specificity at selected cut-off points of the study variables is shown in Table 5, and a comparison of the sensitivity and specificity at selected cut-off points in subgroups according to total PSA value is shown in Table 6. Table 7 shows serum hK11 concentration classified by tumor grade, stage of disease, and Gleason score.

A positive correlation was found between hK11 and patient age in the group of patients with benign prostatic hyperplasia (Spearman correlation coefficient  $r_s=0.242$ ,  $p<0.001$ ) as well as in the group of patients with prostate cancer (Spearman correlation coefficient  $r_s=0.208$ ,  $p=0.009$ ).

The distribution of hK11/total PSA ratio was significantly lower in CaP patients than in BPH patients ( $p<0.001$ ) (Figure 6).

The discriminatory value of the hK11/total PSA ratio was investigated in relation to the total PSA and % free PSA values, by ROC curve analysis (Figure 5). The hK11/total PSA ratio had about the same discriminatory potential as Total PSA.

The hK11/total PSA ratio was analyzed in the subgroup of patients with total PSA = 4-10  $\mu\text{g/L}$ , total PSA >10  $\mu\text{g/L}$ , total PSA < 4  $\mu\text{g/L}$  and %free PSA < 17.8 (Figures 7 to 10). The ratio of hK11/total PSA was also shown to differentiate BPH from prostate cancer in a population of patients with total PSA less than 4ng/ml (Figure 9). In the range of 4-10ng/ml total PSA, the combination of kallikrein 11, total PSA, and % free PSA was especially useful for differentiating BPH from prostate cancer.

The present invention is not to be limited in scope by the specific embodiments described herein, since such embodiments are intended as but single illustrations of one aspect of the invention and any functionally equivalent embodiments are within the scope of this invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

All publications, patents and patent applications referred to herein are incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety. All publications, patents and patent applications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing the domains, cell lines, vectors, methodologies etc. which are reported therein which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

It must be noted that as used herein and in the appended claims, the singular forms "a", "an", and

- 27 -

"the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "a host cell" includes a plurality of such host cells, reference to the "antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Below full citations are set out for the references referred to in the specification.

- 28 -

**Table 1**  
**Descriptive statistics of various variables in serum of 64 BPH patients**

Variables	Mean $\pm$ SE <sup>a</sup>	Range	Percentiles				
			10	25	50 (median)	75	90
hK11 ( $\mu$ g/L)	0.41 $\pm$ 0.05	0.00-2.33	0.06	0.11	0.24	0.62	0.95
Total PSA ( $\mu$ g/L)	5.61 $\pm$ 0.63	0.17 - 26.2	1.05	1.63	4.17	8.02	12.95
Free PSA ( $\mu$ g/L)	1.22 $\pm$ 0.26	0.04 - 15.0	0.22	0.39	0.64	1.41	2.01
% Free/Total PSA	22.3 $\pm$ 0.156	0.85 - 64.0	10.0	13.4	18.5	28.2	38.4
hK11/tPSA	0.14 $\pm$ 0.03	0.00 - 1.28	0.013	0.029	0.063	0.13	0.39

<sup>a</sup> Standard error

5

**Table 2**  
**Descriptive statistics of various variables in serum of 86 CaP patients**

Variables	Mean $\pm$ SE <sup>a</sup>	Range	Percentiles				
			10	25	50 (median)	75	90
hK11 ( $\mu$ g/L)	0.15 $\pm$ 0.02	0.00-0.72	0.00	0.05	0.08	0.25	0.38
Total PSA ( $\mu$ g/L)	9.34 $\pm$ 0.82	0.35 - 48.0	2.86	4.79	7.23	11.50	18.72
Free PSA ( $\mu$ g/L)	1.03 $\pm$ 0.26	0.10 - 23.0	0.27	0.42	0.61	0.97	1.55
% Free/Total PSA	10.9 $\pm$ 0.8	1.9 - 47.9	3.9	6.6	8.6	13.3	21.7
hK11/tPSA	0.028 $\pm$ 0.005	0.00 - 0.31	0.00	0.005	0.014	0.034	0.061

<sup>a</sup> Standard error

- 29 -

Table 3

Descriptive statistics of various variables in serum of BPH and CaP patients

Variables	Mean $\pm$ SE <sup>a</sup>	Range	Percentiles				
			10	25	50 (median)	75	90
BPH (N=247)							
HK11 ( $\mu\text{g/L}$ )	0.19 $\pm$ 0.008	0.010 <sup>b</sup> -1.12	0.062	0.11	0.17	0.22	0.32
Total PSA ( $\mu\text{g/L}$ )	5.37 $\pm$ 0.42	0.58 – 82.9	1.18	1.90	4.00	6.50	11.62
% Free/Total PSA	17.8 $\pm$ 0.58	2.1 - 80.0	6.1	11.7	17.1	22.3	27.8
% hK11/tPSA	6.53 $\pm$ 0.43	0.085 -51.14	0.84	2.03	4.44	8.80	15.23
CaP (N=157)							
HK11 ( $\mu\text{g/L}$ )	0.14 $\pm$ 0.006	0.010 <sup>b</sup> -0.46	0.056	0.091	0.14	0.18	0.24
Total PSA ( $\mu\text{g/L}$ )	13.24 $\pm$ 1.07	0.61- 90.7	3.85	5.72	9.50	14.87	25.25
% Free/Total PSA	9.87 $\pm$ 0.53	1.90 - 56.6	4.05	56.62	8.35	11.87	17.90
% hK11/tPSA	1.90 $\pm$ 0.14	0.02 – 9.38	0.31	0.73	1.30	2.32	4.33

<sup>a</sup> Standard error<sup>b</sup> Half of the assay detection limit

- 30 -

Table 4

Logistic regression analysis of BPH and CaP patients for predicting the presence of prostate cancer.

5

Covariate	Univariate Analysis		Multivariate analysis	
	Crude Odds Ratio	p-value*	Crude Odds Ratio	p-value*
Total PSA	1.17	<0.001	1.05	0.031
% Free/Total PSA Ratio	0.85	<0.001	0.87	0.001
% hK11/Total PSA Ratio	0.66	<0.001	0.76	<0.001

\*Test for trend

Table 5

Comparison of sensitivity and specificity at selected cut-off points of the study variables

Parameter	Cut-off	Specificity (%)	Sensitivity (%)
<u>All Patients (N=404)</u>			
Total PSA	14.92	95	25
	11.62	90	40
	3.86	49	90
	2.89	38	95
% Free/Total PSA	5.95	95	30
	8.35	90	51
	17.8	46	90
	19.2	39	95
% hK11/ Total PSA	0.49	95	19
	0.93	90	33
	3.99	54	90
	5.97	38	95
Ratio Combination <sup>a</sup>	1.50	95	54
	1.82	90	67
	3.19	56	90
	3.69	46	95

<sup>a</sup> Ratio Combination function =  $0.142 * \% \text{ Free/Total PSA} + 0.213 * \% \text{ hK11/Total PSA}$  as it was formulated from the logistic regression analysis.

10



- 31 -

**Table 6**  
**Comparison of sensitivity and specificity at selected cut-off points**  
**in subgroups according to total PSA value**

Parameter	Cut-off	Specificity (%)	Sensitivity (%)
<b><u>Total PSA &lt;4µg/L (N=139)</u></b>			
% Free/Total PSA	7.05	95	6
	8.85	90	18
	19.5	45	90
	20.0	44	95
% hK11/ Total PSA	0.73	95	6
	2.92	90	35
	7.52	57	90
	9.19	43	95
Ratio Combination <sup>a</sup>	2.12	95	18
	2.53	90	31
	4.12	58	90
	4.31	56	95
<b><u>Total PSA 4-10µg/L (N=158)</u></b>			
% Free/Total PSA	5.55	95	25
	8.35	90	53
	15.1	52	90
	18.2	42	95
% hK11/ Total PSA	0.79	95	10
	1.25	90	22
	3.99	28	90
	5.24	16	95
Ratio Combination <sup>a</sup>	1.61	95	47
	1.77	90	58
	2.71	56	90
	3.13	48	95
<b><u>Total PSA &gt;10µg/L (N=107)</u></b>			
% Free/Total PSA	4.65	95	20
	6.35	90	40
	14.6	39	90
	25.8	10	95
% hK11/ Total PSA	0.22	95	13
	0.30	90	15
	1.44	29	90
	1.66	26	95
Ratio Combination <sup>a</sup>	0.83	95	20
	1.33	90	57
	2.22	42	90
	2.88	26	95

<sup>a</sup> Ratio Combination function =  $0.142 * \% \text{ Free/Total PSA} + 0.213 * \% \text{ hK11/ Total PSA}$   
as it was formulated from the logistic regression analysis.

- 32 -

**Table 7**  
**Serum hk11 concentration classified by tumor grade,**  
**stage of the disease and Gleason score**

	Total	Mean <sup>a</sup>	Standard Error <sup>a</sup>	Median <sup>a</sup>	pvalue <sup>b</sup>
<b>Grade</b>					
G1	8	0.171	0.016	0.166	0.036
G2	90	0.152	0.009	0.145	
G3	48	0.120	0.008	0.104	
<b>Stage</b>					
I	13	0.176	0.021	0.174	0.038
II	88	0.141	0.078	0.133	
III/IV	41	0.120	0.011	0.110	
<b>Gleason score</b>					
≤ 6	55	0.151	0.117	0.146	0.176
>6	72	0.131	0.008	0.125	

<sup>a</sup> ug/L<sup>b</sup> Calculated by the Mann-Whitney U Test

## FULL CITATIONS FOR REFERENCES REFERRED TO IN THE SPECIFICATION

1. Oesterling JE: Prostate specific antigen: a critical assessment of the most useful tumor marker for  
5 adenocarcinoma of the prostate. *J Urol* 145: 907-923, 1991.
2. Barry MJ: Clinical practice. Prostate-specific-antigen testing for early diagnosis of prostate cancer.  
*N Engl J Med* 344: 1373-1377, 2001
3. Oesterling JE, Jacobsen SJ, Klee GG, et al: Free, complexed and total serum prostate specific  
antigen: the establishment of appropriate reference ranges for their concentrations and ratios. *J Urol*  
10 154: 1090-1095, 1995.
4. Stephan C, Jung K, Lein M, et al: Molecular forms of prostate-specific antigen and human  
kallikrein 2 as promising tools for early diagnosis of prostate cancer. *Cancer Epidemiol Biomarkers  
Prev* 9: 1133-1147, 2000.
5. Catalona WJ, Partin AW, Slawin KM, et al: Use of the percentage of free prostate-specific antigen  
15 to enhance differentiation of prostate cancer from benign prostatic disease: a prospective  
multicenter clinical trial. *JAMA* 279: 1542-1547, 1998.
6. Yoshida S, Taniguchi M, Suemoto T, et al: cDNA cloning and expression of a novel serine  
protease, TLSP. *Biochim Biophys Acta* 1399: 225-228, 1998.
7. Mitsui S, Yamada T, Okui A, et al: A novel isoform of a kallikrein-like protease, TLSP/hippostasin,  
20 (PRSS20), is expressed in the human brain and prostate. *Biochem Biophys Res Commun* 272: 205-  
211, 2000.
8. Nakamura T, Mitsui S, Okui A, et al: Alternative splicing isoforms of hippostasin  
(PRSS20/KLK11) in prostate cancer cell lines. *Prostate* 49: 72-78, 2001.
9. Yousef GM, Scorilas A, and Diamandis EP: Genomic organization, mapping, tissue expression, and  
25 hormonal regulation of trypsin-like serine protease (TLSP PRSS20), a new member of the human  
kallikrein gene family. *Genomics* 63: 88-96, 2000.
10. Diamandis EP, Yousef GM, Clements J, et al: New nomenclature for the human tissue kallikrein  
gene family. *Clin Chem* 46: 1855-1858, 2000.
11. Diamandis EP, Okui A, Mitsui S, et al: Human Kallikrein 11: A new biomarker of prostate and  
30 ovarian carcinoma. *Cancer Res* 62: 295-300, 2002.
12. Catalona WJ, Partin AW, Finlay JA, et al: Use of percentage of free prostate-specific antigen to  
identify men at high risk of prostate cancer when PSA levels are 2.51 to 4 ng/mL and digital rectal  
examination is not suspicious for prostate cancer: an alternative model. *Urology* 54: 220-224, 1999.
13. Jung K, Elgeti U, Lein M, et al: Ratio of free or complexed prostate-specific antigen (PSA) to total  
35 PSA: which ratio improves differentiation between benign prostatic hyperplasia and prostate  
cancer? *Clin Chem* 46: 55-62, 2000.
14. Oesterling JE, Jacobsen SJ, and Cooner WH: The use of age-specific reference ranges for serum  
prostate specific antigen in men 60 years old or older. *J Urol* 153: 1160-1163, 1995.
15. Magklara A, Scorilas A, Catalona WJ, et al: The combination of human glandular kallikrein and

- 34 -

free prostate- specific antigen (PSA) enhances discrimination between prostate cancer and benign prostatic hyperplasia in patients with moderately increased total PSA. Clin Chem 45: 1960-1966, 1999.

- 35 -

**I Claim:**

1. A method for detecting prostate cancer in a subject comprising measuring kallikrein 11 and prostate specific antigen (PSA) in a sample from the subject.
2. A method for detecting prostate cancer in a subject comprising:
  - 5 (a) determining the amount of kallikrein 11 in a sample from the subject;
  - (b) determining the amount of PSA in the sample;
  - (c) mathematically combining the results of step (a) and step (b); and
  - (d) relating the combination to the presence of prostate cancer.
3. A method as claimed in claim 1 or 2 wherein in step (a) kallikrein 11 is determined using antibodies that specifically bind to kallikrein or a part thereof.
- 10 4. A method as claimed in claim 1, 2, or 3 wherein in step (b) the PSA is measured using antibodies that specifically bind to PSA or a part thereof
5. A method as claimed in claim 2 wherein in step (a) kallikrein 11 is determined in the sample by
  - 15 (a) incubating a sample from the subject with a first antibody that specifically binds kallikrein 11 which is directly or indirectly labeled with a detectable substance, and a second antibody that specifically binds kallikrein 11 which is immobilized; and
  - (b) detecting the detectable substance thereby determining kallikrein 11 in the sample.
6. A method as claimed in claim 2, 3, 4, or 5 wherein in step (b) total PSA is determined in the sample.
- 20 7. A method as claimed in any of claims 2 to 6 wherein the combination is a ratio of kallikrein 11 to total PSA, or the inverse thereof.
8. A method as claimed in any preceding claim which further comprises the step of determining the % free PSA and relating the combination and % free PSA to the presence of prostate cancer.
9. A method as claimed in any preceding claim wherein the combination is compared to a predetermined standard.
- 25 10. A method for distinguishing prostate cancer from benign prostatic hyperplasia (BPH) in a subject comprising determining the amount of kallikrein 11 contained in a sample from the subject, and relating the amount to the presence of prostate cancer or BPH in the subject.
11. A method as claimed in claim 10 wherein the kallikrein 11 is measured using antibodies that specifically bind to kallikrein 11 or a part thereof.
- 30 12. A method as claimed in claim 10 or 11 wherein the amount of kallikrein 11 in the sample is compared to an amount determined for a standard.
13. A method as claimed in claim 12 wherein the standard is an amount of kallikrein 11 associated with prostate cancer.
- 35 14. A method as claimed in claim 13 wherein an amount of kallikrein 11 in the sample greater than the standard is indicative of BPH.
15. A method as claimed in claim 12 wherein the standard is an amount of kallikrein 11 associated with BPH.
16. A method as claimed in claim 15 wherein an amount of kallikrein 11 in the sample lower than the

- 36 -

standard is indicative of prostate cancer.

17. A method for distinguishing prostate cancer from benign prostatic hyperplasia (BPH) in a subject comprising:
  - (a) determining the amount of kallikrein 11 contained in a sample from the subject;
  - 5 (b) determining the amount of total PSA contained in the sample;
  - (c) mathematically combining the results of (a) and (b);
  - (d) relating the combination to the presence of BPH or prostate cancer.
18. A method as claimed in claim 17 wherein the combination is a ratio of kallikrein 11 to total PSA, or the inverse thereof.
- 10 19. A method as claimed in claim 17 or 18 wherein the kallikrein 11 is measured using antibodies specifically reactive with kallikrein 11 or a part thereof.
20. A method as claimed in claim 17, 18 or 19 wherein the PSA is measured using antibodies specifically reactive with PSA or a part thereof.
21. A method as claimed in any one of claims 17 to 20 wherein the combination is compared to a combination for a standard.
- 15 22. A method as claimed in claim 21 wherein the standard is a combination associated with prostate cancer.
23. A method as claimed in claim 22 wherein a combination in the sample is greater than the standard is indicative of BPH.
- 20 24. A method as claimed in claim 21 wherein the standard is a combination associated with BPH.
25. A method as claimed in claim 24 wherein a combination in the sample lower than the standard is indicative of prostate cancer.
26. A method of any one of claims 17 to 25 further comprising determining the percentage of free PSA and correlating the percentage free PSA and the combination to the presence of prostate cancer or BPH in the subject.
- 25 27. A method for determining the presence of BPH or prostate cancer in a subject comprising:
  - (a) providing a first binding agent that specifically binds to kallikrein 11;
  - (b) providing a second binding agent that specifically binds to PSA;
  - (c) contacting the first agent and second agent with the sample under a condition that allows the formation of a first complex comprising the first agent and the kallikrein 11, and a second complex comprising the second agent and the PSA;
  - 30 (d) determining the presence or amount of the first and second complexes;
  - (e) mathematically combining the amount of the first and second complexes; and
  - (f) relating the combination to the presence of BPH or prostate cancer.
- 35 28. A method as claimed in claim 27 wherein the combination is a ratio of the first complex to the second complex, or the inverse thereof.
29. A method of claim 27 or 28 wherein the binding agents are antibodies.
30. A method of claim 27 or 28 further comprising determining the percentage of free PSA and correlating the percentage free PSA and the combination to the presence of prostate cancer or BPH

in the subject.

31. A method of any preceding claim wherein the subject has total PSA between about 4-10 ng/ml.
32. A method of any preceding claim wherein the subject has total PSA less than 4 ng/ml.
33. A method of any preceding claim wherein the sample is a mammalian tissue sample.
- 5 34. A method of any preceding claim wherein the sample is a sample of human physiological fluid.
35. A method of any preceding claim wherein the sample is serum, seminal plasma, urine, or plasma.
36. A method of improving the accuracy of a diagnosis of prostate cancer comprising the steps of: a) performing a method of any of the preceding claims; and b) performing at least one of a test for free PSA and a digital rectal examination.
- 10 37. A method for screening prostate cancer by determining the ratio between kallikrein 11:total PSA in a subject's serum.
38. A method for differentiation between BPH or prostate cancer by determining the ratio between kallikrein 11:total PSA in a subject's serum.
39. A kit for determining the presence of BPH or prostate cancer in a subject, comprising a known  
15 amount of a binding agent that specifically binds to kallikrein 11 wherein the binding agent comprises a detectable substance, or it binds directly or indirectly to a detectable substance.
40. A kit for determining the presence of BPH or prostate cancer in a subject, comprising:  
(a) a known amount of a first binding agent that specifically binds to kallikrein 11; and  
(b) a known amount of a second binding agent that specifically binds to PSA;  
20 wherein the first and second binding agents comprise a detectable substance, or bind directly or indirectly to a detectable label.
41. A method for screening a subject for prostate cancer comprising (a) obtaining a biological sample from a subject; (b) detecting the amount of kallikrein 11 and PSA in said sample; and (c) comparing said amount of kallikrein 11 and PSA or a mathematical combination thereof to a predetermined  
25 standard, where detection of a significant difference in kallikrein 11 and PSA or a mathematical combination thereof compared to a standard is indicative of prostate cancer.
42. A method for monitoring the progression of prostate cancer in a patient, the method comprising: (a) detecting in a sample from the patient at a first time point, kallikrein 11 and PSA (b) repeating step (a) at a subsequent point in time; and (c) comparing levels detected in steps (a) and (b) or a  
30 mathematical combination thereof, and thereby monitoring the progression of prostate cancer.
43. A method for assessing the potential efficacy of a test agent for inhibiting prostate cancer in a patient, the method comprising comparing: (a) levels of kallikrein 11 and PSA or a mathematical combination thereof in a first sample obtained from a patient and exposed to the test agent; and (b) levels of kallikrein 11 and PSA or a mathematical combination thereof in a second sample obtained  
35 from the patient, wherein the sample is not exposed to the test agent, wherein a significant difference in kallikrein 11 and PSA or a mathematical combination thereof in the first sample, relative to the second sample, is an indication that the test agent is potentially efficacious for inhibiting prostate cancer in the patient.
44. A method of assessing the efficacy of a therapy for inhibiting prostate cancer in a patient, the

- 38 -

method comprising comparing: (a) levels of kallikrein 11 and PSA or a mathematical combination thereof in a first sample obtained from the patient, and (b) levels of kallikrein 11 and PSA or a mathematical combination thereof in a second sample obtained from the patient following therapy, wherein a significant difference in kallikrein 11 and PSA or a mathematical combination thereof in the second sample, relative to the first sample, is an indication that the therapy is efficacious for inhibiting prostate cancer in the patient.

45. A method of selecting an agent for inhibiting prostate cancer in a patient the method comprising (a) obtaining a sample of cells affected by breast or ovarian cancer from the patient; (b) separately exposing aliquots of the sample in the presence of a plurality of test agents; (c) comparing levels of kallikrein 11 and PSA or a mathematical combination thereof in each of the aliquots; and (d) selecting one of the test agents which alters kallikrein 11 and PSA or a mathematical combination thereof in the aliquot containing that test agent, relative to other test agents.

46. A method of inhibiting prostate cancer in a patient, the method comprising (a) obtaining a sample comprising cells affected by prostate cancer from the patient; (b) separately maintaining aliquots of the sample in the presence of a plurality of test agents; (c) comparing levels of kallikrein 11 and PSA or a mathematical combination thereof in each of the aliquots; and (d) administering to the patient at least one of the test agents which alters kallikrein 11 and PSA or a mathematical combination thereof in the aliquot containing that test agent, relative to other test agents.

47. A method of assessing the potential of a test compound to contribute to prostate cancer, the method comprising: (a) maintaining separate aliquots of cells affected by the breast or ovarian cancer in the presence and absence of the test compound; and (b) comparing levels of kallikrein 11 and PSA or a mathematical combination thereof in each of the aliquots, and wherein a significant difference in kallikrein 11 and PSA or a mathematical combination thereof in the aliquot maintained in the presence of the test compound, relative to the aliquot maintained in the absence of the test compound, is an indication that the test compound possesses potential compound to contribute to prostate cancer.

48. A kit for carrying out a method of any preceding claim.


49. A method of conducting a drug discovery business comprising:

- (a) providing a method as claimed in claim 45 for selecting an agent that inhibits prostate cancer in a patient;
- (b) conducting therapeutic profiling of agents identified in step (a), or further analogs thereof, for efficacy and toxicity in animals; and
- (c) formulating a pharmaceutical preparation including one or more agents identified in step (b) as having an acceptable therapeutic profile.




1/10

Figure 1




2/10

Figure 2




3/10

Figure 3




4/10

Figure 4




5/10

Figure 5




6/10

Figure 6



7/10

Figure 7




SUBSTITUTE SHEET (RULE 26)

SUBSTITUTE SHEET (RULE 26)

8/10


Figure 8






9/10

Figure 9



10/10

Figure 10



- 1 -

## SEQUENCE LISTING

<110> Mount Sinai Hospital  
 5 <120> Methods of Detecting Prostate Cancer  
 <130> T01114-0017-WO  
 <140>  
 10 <141>  
 <150> US 60/414,314  
 <151> 2002-09-26  
 15 <160> 3  
 <170> PatentIn version 3.2  
 <210> 1  
 20 <211> 282  
 <212> PRT  
 <213> homo sapiens  
 <300>  
 25 <308> Genbank/AB012917  
 <309> 1998-04-10  
 <313> (1)..(282)  
 <400> 1  
 30 Met Gln Arg Leu Arg Trp Leu Arg Asp Trp Lys Ser Ser Gly Arg Gly  
 1 5 10 15  
 35 Leu Thr Ala Ala Lys Glu Pro Gly Ala Arg Ser Ser Pro Leu Gln Ala  
 20 25 30  
 40 Met Arg Ile Leu Gln Leu Ile Leu Leu Ala Leu Ala Thr Gly Leu Val  
 35 40 45  
 45 Gly Gly Glu Thr Arg Ile Ile Lys Gly Phe Glu Cys Lys Pro His Ser  
 50 55 60  
 Gln Pro Trp Gln Ala Ala Leu Phe Glu Lys Thr Arg Leu Leu Cys Gly  
 65 70 75 80  
 50 Ala Thr Leu Ile Ala Pro Arg Trp Leu Leu Thr Ala Ala His Cys Leu  
 85 90 95  
 55 Lys Pro Arg Tyr Ile Val His Leu Gly Gln His Asn Leu Gln Lys Glu  
 100 105 110

- 2 -

Glu Gly Cys Glu Gln Thr Arg Thr Ala Thr Glu Ser Phe Pro His Pro  
 115 120 125

5 Gly Phe Asn Asn Ser Leu Pro Asn Lys Asp His Arg Asn Asp Ile Met  
 130 135 140

10 Leu Val Lys Met Ala Ser Pro Val Ser Ile Thr Trp Ala Val Arg Pro  
 145 150 155 160

15 Leu Thr Leu Ser Ser Arg Cys Val Thr Ala Gly Thr Ser Cys Leu Ile  
 165 170 175

Ser Gly Trp Gly Ser Thr Ser Ser Pro Gln Leu Arg Leu Pro His Thr  
 180 185 190

20 Leu Arg Cys Ala Asn Ile Thr Ile Ile Glu His Gln Lys Cys Glu Asn  
 195 200 205

25 Ala Tyr Pro Gly Asn Ile Thr Asp Thr Met Val Cys Ala Ser Val Gln  
 210 215 220

30 Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Val  
 225 230 235 240

35 Cys Asn Gln Ser Leu Gln Gly Ile Ile Ser Trp Gly Gln Asp Pro Cys  
 245 250 255

Ala Ile Thr Arg Lys Pro Gly Val Tyr Thr Lys Val Cys Lys Tyr Val  
 260 265 270

40 Asp Trp Ile Gln Glu Thr Met Lys Asn Asn  
 275 280

45 <210> 2  
 <211> 9120  
 <212> DNA  
 <213> homo sapiens

50 <300>  
 <308> Genbank/AF164623.  
 <309> 1999-07-01  
 <313> (1)..(9120)

55 <400> 2  
 tgataatagt gttctctctc ctcattgggc agggccccag ccattgtcct tgagagaatg 60  
 ctgcactctt tatgttgtct tgacagcctc cctgagatt ggtcattaat gactgtgctc 120

- 3 -

	tctctctca ttggtcaggg cccagccat tgtccttgag agaacctctg tcctttatgg	180
	agttccaccc ttcttcctg ggattggccc ctagagacag tggttcttct cttttggtta	240
5	gccattgcca ttgtcctccg ggaaagtgat tatactcttt tgtctaataa ccagacttgg	300
	agccctcccc aaggcccagg actgggttga agggttgggg aggaaaacag aaataagatg	360
10	tctcccttgt tcagacagta cttctcttcc cttccagggt gattctgggg gccccctgg	420
	gtgtggggga gtccttcaag gtctggtgtc ctgggggtct gtggggccct gtggacaaga	480
	tggcatccct ggagctctaca cctatatatt caagtatgtg gactggatcc ggatgatcat	540
15	gaggaacaac tgacctgttt cctccacctc cccccacc ccttaacttg ggtaccctc	600
	tggccctcag agcaccaata tctctccat cacttcccct agtccactc ttgttggcct	660
20	gggaacttct tggaacttta actcctgcca gcccttctaa gaccacagag cggggtgaga	720
	gaagtgtgca atagtctgga ataaatataa atgaaggagg ggccatgtct gtccatttga	780
	agtcctcatg ctggttgaga ctggaagaag gactcagcag tttccctatc tcataggagt	840
25	agaaacagag ctcaaataag gccaggcaca gtggctcaca cctgtaatcc catcactttg	900
	ggaagctgag gcagggtgat cacctgaggt caggaactcg ggaccagcct ggtcaacata	960
30	gtgaaacccc aactctacta aaaatgcaaa aattagccag gcatggtggc gcatgcctgt	1020
	aatcccagct actcaggagg ctgagacagg agaatagcat gaaccctga ggcagaggct	1080
	gcagcgagcc gagattgaac cattacactc cagcctgggc gacagagcga gactccatct	1140
35	caaaaacaaa caaacaacaaa acccagtgtc caaataggat gaggtcttcc cctgagtagt	1200
	tactcagaaa tggagtagaa aaagttactt ttaataatat aggccgggtg cagtggccca	1260
40	cgctgtaat ccagcactt tgggaggccg aggtgggagg atggcttgag ctgagatttc	1320
	gagatcagcc tggcaacaca gtgaaatctt gtcactacaa aaacacaaaa aattagctgg	1380
	gtgtggtggt gcgtgcctgt agtcccagct acttgggaag ctgaggtggg aggatcacc	1440
45	gagccgggga ggtggaggct gcaaagagcc gagatcatgc cactgcactc cagcctgggc	1500
	aataaagtga gacottgtct caaaaacaaa aaccagcaa tataaataag acacatgttt	1560
50	cttcatctgg cataatagaa atagtccca gagcttataa gcttttcaag agtcacaaaa	1620
	agacccgaaa aagaaaaaga aaattgttag ctccaaaata ccagatgaaa gctgcaaagt	1680
	caacatttat gaccatttaa tccaatgtcc ataaaacgta gcattcttcc cactagccaa	1740
55	ctgcagttta ctttcttgta atgaagcata cattgtatct ttaatgtggg acgtggcttt	1800
	gttctaataa gacgaagggt ggagtgcagg cttggaaagc aggagagctc agcctacgtc	1860
	tttaatctc ctgcccacc cttggattct gtctccactg ggactcaaga ggtgaggaga	1920

- 4 -

	gaccatctcc ccaaatacac tgaagggaaa ctggaggagg gagggagtga ggggtgatca	1980
	taccagcggg ggcacatttg ctgagcccc ccgcagtctg ctctttccaa gtggaccctc	2040
5	ctggaagcct gatcccaacc tcccctgcaa gcaggtctgt ccccccatc tctcagatga	2100
	agaaactgag ccttgcaagg gtggagtccc ttgtccccac gtcataaggg tagtcatagt	2160
10	agtaggaaga ggaagcacct aggtttgagg ccagggtctg ctgctgtcag aacctaggcc	2220
	ctcccctgcc ttgtccaca cctggtcagg ggagagaggg gaggaagcc aagggaagg	2280
	acctaaactga aaacaacaa gctgggagaa gcaggaatct gcgctcgggt tccgcagatg	2340
15	cagaggttga ggtggctgcg ggactggaag tcatcgggca gaggtctcac agcagccagt	2400
	aagtgaacag ctggactcgg gctgcctggg cggcaggag aagcgggag gggaagggtc	2460
20	agcagaggag cgaggcccca gaggagccct ggggtggagc acagccaagg gctctgttcc	2520
	ctttcctgga ctcggttcc acaggccctg acctgcctcc cccaccctcc ggtcctgccc	2580
	ctgtgcctgg cagcagcccc acctgtgtga catcccagca cccccccct ctocctgcaa	2640
25	aggagaaggg agcggcctag gggaggccag gggccacct gggctggggc tgtggagagg	2700
	gagtggctgg gacgggagga aaaagagaga cggagattag atggaagaag agggatttca	2760
30	agacaaattg ccagagatgc agtcagagag actgactgag agacacaaag atagaaggaa	2820
	ttagagaaaag ggccacacag agccagacag agagagaaga gtggagatgg agacagggac	2880
	gaggacagag aaaggcagac agacacatag ggacagaaag agaaaaatca cacaaagtca	2940
35	gaattactga atgacaggga atgacacata gaacgagaca cagattcaga gactcagggc	3000
	agggaaagga aggtgcaga cagacagaca gacagaggga ggctgagaca cagggagaag	3060
40	aggggcttgg agaggtggca caggcaggca gccagtgcct cagaggcctc cggggagggc	3120
	cctcacacac accccgcccc ggggcattaa ggcagggtt ggaggccagt catcctgggc	3180
	cggcccaggg ccgccccct gccagccgc ctgcctggtg cctggcacct ggcgtccaa	3240
45	cccagcctac ctgctgtagc tgccgccact gccgtctccg ccgccactgg gccccagag	3300
	cccagcccc agagcctgtg agtccaggag gaaagggaag ctgcccctcc ccgtccaggt	3360
50	gtcagccctc cccaaggaca cctgtccac tggggacccc atttctccct ctgctctgtc	3420
	cttctctgt tgggtgggg ttctggcct ctctctacac ctctcacctc cgatggctgt	3480
	ccgcagcctc agttacctt aatctccatg gcttcagctg ctgagctggc cctctgtctc	3540
55	cacccccgt ggccaggga cggaggga ctggccctcc cctcgaccag ccccgccag	3600
	ctttgcttgg ctgtccttca aaagggcagg ggtttggcg acagggttc agcaagccgg	3660

- 5 -

	gtgatggggg tcccagacat tgtctggggc tgagccccct actcccctcc agcagacctc	3720
	aaaggtcca tatcgctctg ctgcgaagac aatgaaaaag gggtggtac ggaacggtgt	3780
5	ctggttcccc ttgtccttcc accccaagct gctggggcct ggccagctct caaggcaaga	3840
	aggaaaacat cctctgacat gtgccgggga ggtcccatgg ctgacttgaa cagggccgaa	3900
10	ccatggcttg acagctcaaa gccctccca acgacttcca catggttctt ggtatctcgg	3960
	aagcttctag ctgtgaccag gccctctcca aggccaccct agacacctaa gatataTTTT	4020
	aagtgtttgg agatctgagt gctgtgagaa acaggggatt tccccaacct tgtttctccc	4080
15	aagtggggag cgggagcagg tgaggagag aggagagggc atgagccagc cccccctcc	4140
	cgatttcccc gtaaagtgat gcggcccat gtccctcctt gttcccagag gaacctgggg	4200
20	cccgctcctc cccctccag gccatgagga ttctgcagtt aatcctgctt gctctggcaa	4260
	caggtagca ggggatggg gcagggcagg atcctccctc ttgaatctct gggatccct	4320
	aacctctgt gtctggacag tgacagggt gattccaaat tacagaacaa ccataaggc	4380
25	acctgaactg gagcagtgg catgagggcc tggatgccct tctagataat ccttttaaat	4440
	gccaaaggag gagaggtcaa gggggtcgta aagggtcccg tggaggggt gaggaagtgt	4500
30	gagttggggg agcagtcact caaagcgccc aggacaggg ctactgacca accagtatgg	4560
	aagtatttcc tttttttttt ttcccagaga caaagtcttg ctctattgtc caggctggag	4620
	tgccgtggtg ccaacacggc tcaactgcagt cttgacttcc cgggcttaag tgatccttaa	4680
35	gccatctcag cttccccggt agctgggacc acaggcacct gccaccaagc caggctaatt	4740
	gtttaattgt ttgtagagat ggggagggag gtctcactat gtttgcttag gctgatctag	4800
40	aactcctggg ctcaagtaat cctccacct tagcctctca aagtgtggg attacaggca	4860
	tgagccactg ctttgacct tatggaagta ttttcactct ttaatacccg accccagcat	4920
	ccagggaac ccagagggac accagaccag ggcccagacc accactctc tttctctcct	4980
45	ccccaccccc atttctggga gtccctctgg tctaccacct ctccttcttg agccccctct	5040
	tttgcctca cccctccag ggcttgtagg gggagagacc aggatcatca aggggttcga	5100
50	gtgcaagcct cactcccagc cctggcaggc agccctgttc gagaagacgc ggctactctg	5160
	tggggcgacg ctcatcgccc ccagatggct cctgacagca gccactgcc tcaagccgtg	5220
	ggtgcggggg ctggggcggt gccgggggtg ggggctggga atggggagat ggatggagag	5280
55	aagctcaggg ataggggtgc tggttaaggg attagagatg gggatgggta gtgtcagcaa	5340
	ggttgatggg ctcgagtgg tattgaaggt ggggggatga atgggggttg gatggggcta	5400
	tggctgggaa gggggcttcg gtgggagacg tggaagaggt tggaagcaga gcgatgtttc	5460

	ttcatcctca aaggtgtcac tcacctctcc caccatgtc tcccccgacc tttcctcctc	5520
5	caactactgt ctctcccacc tcagccgcta catagttcac ctggggcagc acaacctcca	5580
	gaaggaggag ggctgtgagc agaccggac agccactgag tccttcccc accccggctt	5640
	caacaacagc ctccccaaca aagaccaccg caatgacatc atgctggtga agatggcatc	5700
10	gccagtctcc atcacctggg ctgtgcgacc cctcaccctc tcctcacgct gtgtcactgc	5760
	tggcaccagc tgcctcattt ccggctgggg cagcacgtcc agccccagt gtaggagcac	5820
15	cagaggggaa cctggcaggg ggtggtgagg agggagtggg caggattgtg gaagggttca	5880
	gggcatcaga gatgcggttc acagtgcga tgtgggataa gttgagagga tgtgtggaaa	5940
	acgtcaggat aggggggtgg ggacaaaagt tggggccttg gagtcagacg gacgggatat	6000
20	gcaatcatac atccataacc tcctggttgt aagaccttag gcaagcagct tcacctctc	6060
	gaatcttgat tttcttctct ataaaatgag aatgattata cccacctgtc aggattggat	6120
25	tagagataat gtatatcaag caactgacat aaatcattta ttggatagca ggctgggcac	6180
	cgtggctcac gcctgtaatc ccagcacttt gggaggccga ggtgggaaga tcacctgagg	6240
	tcaggacttt gataccagcc tggccaacgt ggtgaaatcc catctctact aaaaatgtga	6300
30	aaattagttg ggcgtggttg tgtgcgcctg taatcccagc tactcgggag gttgaggcag	6360
	gagaatcgct taaacttggg agacggagggt tgcagtgagc caagatcacg ccactgcact	6420
35	ccagcctggg caacagagca agactctgtc tcgaaaaaa aaaaaaaaaa gctggatagc	6480
	attgtgtgtg ctattgttac aagaagagag gtgagttggc tgcgtctaag gacagggatt	6540
	ccccagggg cgggatcaca gcaagcactg cattagggga ggtggcaggg ggctcattcc	6600
40	cacagccctc cagctgttt ccacagtacg cctgcctcac accttgcat gcgccaacat	6660
	caccatcatt gagcaccaga agtgtgagaa cgcctacccc ggcaacatca cagacaccat	6720
45	ggtgtgtgcc agcgtgcagg aagggggcaa ggactcctgc caggtcagtg tggctctcaa	6780
	ccacagcccc atcccctcc ccagcttcaa tgacatcttt accgacatcc acaatttcat	6840
	ccccaacctc aaccgcoga cccctgcaac tcccaatcca tctcttcccc tgttcccggt	6900
50	tctgacctca gcacaaactt cagctccatc cccgtttcca caccatttcc agctccaacc	6960
	atccccaaac tcgtttttga gcctaaccoc atcctttatc ccaccataa tcccagcttt	7020
55	atcgctaaac ctatcacctt tcccagtgcc taccatcct gtctcggccc cactcctaag	7080
	caccgtcccc acctcctccc tggctaacac catgtcaac gctttctctg accgacattc	7140
	tctctccccg tgcccagggt gactccgggg gccctctggt ctgtaaccag tctcttcaag	7200



- 7 -

	gcattatctc ctggggccag gatccgtgtg cgatcaccgc aaagcctggt gtctacacga	7260
	aagtctgcaa atatgtggac tggatccagg agacgatgaa gaacaattag actggaccca	7320
5	cccaccacag cccatcaccc tccatttcca cttggtgttt ggttcctggt cactctgtta	7380
	ataagaaacc ctaagccaag accctctacg aacattcttt gggcctcctg gactacagga	7440
10	gatgctgtca ctttaataatc aacctggggt tcgaaatcag tgagacctgg attcaaattc	7500
	tgccctgaaa tattgtgact ctgggaatga caacacctgg tttgttctct gttgtatccc	7560
	cagccccaaa gacagctcct ggccatatat caaggtttca ataaatattt gctaaatgag	7620
15	tgaatctact gagtgcttac tatgtgctag accctgatcc aatggctttt attttatttt	7680
	attttttgac agagtctcgc tctgtcaccc aggctggagt acagtgggtc tatctctgct	7740
20	cactgcaacc tccacctcct gggttcaagc aattctcctg cctcagcctc ctgaatagct	7800
	gggattacag gtgcctacca ccacatccgg ctaatttttg tatttttttag tagagatggg	7860
	gcttcaccat gttggccagg ctggtctcga actcctgacc tcagatgac tgccctcctt	7920
25	ggcctcccaa actcctggga ttacagacgt gagccaccgc gcccgccgg ctttcattta	7980
	ttaattaaaa gaaattaaat taattaatct atttaggaga cagtcttget ctgttgccca	8040
30	ggctggagtg cagtaacaat cacagctcac ggcaatctca atttctcctg gtcaagtgat	8100
	tgtctctcct cagcctccag agtagctggg actacaggca catgccagca agcccagcta	8160
	atttttgtat ttttcgtaga gacagaggtc tcagtatgtt gccccggcta gtctcaaact	8220
35	cctgggctca agcagtctgt cctcctcagc ctccaaaagt ggtgagatta caggcatgag	8280
	tcgctgtgcc tggcctccaa gcactttcaa atgtatcaac ttaatcctca caaaaccctg	8340
40	tgaggctcgt actgttttca tacctatttt atagttgaag aaacagacac agagaagcaa	8400
	agtcacttgc tcacagtcac gtggctagga gagcaaggat ctgaagcaag gcgatctctt	8460
	aattaccaag tgatgttctt ggagtaaggc tctgtttgtt tcctttcctg taaaatgctg	8520
45	catgcaaaag tataacacag taagtaaaga agtcagttag cctgcacata ctaagacct	8580
	accaaaggag cttattgttt ttctccaact tccatgatag gtaattagat agtggagacc	8640
50	tctgctggcc aatatggtag ccactaacgc cagctggctc ttccaattaa aattacataa	8700
	agccagaaat gtaactcctc tgtctcactt gttatatctc caaggctgga tagccacatg	8760
	tgactggtgg tggtcggatt agctagtgca tataaaacat cactgcagaa agttcagctg	8820
55	agcagcactg agttagatgg cctctgaaga ggatgtccca cggagagaat ccagaactca	8880
	ggatcttttt tttttttttt ctttgcgaca gagtcttget ctgtcaccca ggctggagtg	8940
	cagtggcgtg atctcggctc actgcaactt ctgcctccca ggttcaagca atttctcctg	9000

- 8 -

ctcagcctcc ctagtagctg ggactacagg cctgtgccaa catccccagc taatTTTTgt 9060  
 gtctTTTTtag tagagatggg gtttactat gttggccagg ctggtctcga actcctgacc 9120  
 5  
 <210> 3  
 <211> 1186  
 <212> DNA  
 10 <213> homo sapiens  
 <400> 3  
 aggaatctgc gctcgggttc cgcagatgca gaggttgagg tggctgcggg actggaagtc 60  
 15 atcggggcaga ggtctcacag cagccaagga acctggggcc cgctcctccc ccctccaggc 120  
 catgaggatt ctgcagttaa tcctgcttgc tctggcaaca gggctttagt ggggagagac 180  
 caggatcatc aaggggttcg agtgcaagcc tactcccag ccctggcagg cagccctgtt 240  
 20 cgagaagacg cggctactct gtggggcgac gtcacatgcc ccagatggc tcctgacagc 300  
 agcccactgc ctcaagcccc gctacatagt tcacctggg cagcacaacc tccagaagga 360  
 25 ggagggtgtg gagcagacc ggacagccac tgagtccttc cccaccccc gcttcaacaa 420  
 cagcctcccc acaaagacc accgcaatga catcatgctg gtgaagatgg catcgccagt 480  
 ctccatcacc tgggtgtgac gaacctcac cctctctca cgctgtgtca ctgctggcac 540  
 30 cagctgcctc atttcgggtt ggggcagcac gtccagcccc cagttacgcc tgcctcacac 600  
 cttgcgatgc gccaacatca ccatcattga gcaccagaag tgtgagaacg cctaccccg 660  
 35 caacatcaca gacacatgg tgtgtgccag cgtgcaggaa gggggcaagg actcctgcca 720  
 ggggtgactcc gggggccctc tggctgttaa ccagtctctt caaggcatta tctcctgggg 780  
 ccaggatccg tgtgcgatca cccgaaagcc tgggtgtctac acgaaagtct gcaaatatgt 840  
 40 ggactggatc caggagacga tgaagaacaa ttagactgga cccaccacc acagcccatc 900  
 accctccatt tccacttggg gtttgggtcc tgttactct gttaataaga aaccctaagc 960  
 45 caagaccctc tgcgaacatt ctttgggcct cctggactac aggagatgct gtcacttaat 1020  
 aatcaacctg gggttcgaaa tcagtgaac ctggattcaa attctgcctt gaaatattgt 1080  
 gactctggga atgacaacac ctggtttgtt ctctgttgta tcccagccc caaagacagc 1140  
 50 tcctggccat atatcaaggt ttcaataaat atttgctaaa tgagtg 1186